

The non-canonical tomato yellow leaf curl virus recombinant that displaced its parental viruses in southern Morocco exhibits a high selective advantage in experimental conditions

Zineb Belabess,^{1,2,3} Michel Peterschmitt,¹ Martine Granier,¹ Abdessalem Tahiri,² Abdelali Blenzar³ and Cica Urbino¹

Correspondence
Cica Urbino
cica.urbino@cirad.fr

¹CIRAD-INRA-SupAgro, UMR BGPI, CIRAD, TA A -54K, Campus International de Baillarguet, F-34398 Montpellier, France

²Ecole Nationale d'Agriculture de Meknès, BPS 40, Meknès, Morocco

³Faculté des Sciences de Meknès BP 11201, Avenue Zitoune, Meknès, Morocco

Recombination events are frequently inferred from the increasing number of sequenced viral genomes, but their impact on natural viral populations has rarely been evidenced. TYLCV-IS76 is a recombinant (*Begomovirus*, *Geminiviridae*) between the Israel strain of tomato yellow leaf curl virus (TYLCV-IL) and the Spanish strain of tomato yellow leaf curl Sardinia virus (TYLCSV-ES) that was generated most probably in the late 1990s in southern Morocco (Souss). Its emergence in the 2000s coincided with the increasing use of resistant tomato cultivars bearing the *Ty-1* gene, and led eventually to the entire displacement of both parental viruses in the Souss. Here, we provide compelling evidence that this viral population shift was associated with selection of TYLCV-IS76 viruses in tomato plants and particularly in *Ty-1*-bearing cultivars. Real-time quantitative PCR (qPCR) monitoring revealed that TYLCV-IS76 DNA accumulation in *Ty-1*-bearing plants was significantly higher than that of representatives of the parental virus species in single infection or competition assays. This advantage of the recombinant in *Ty-1*-bearing plants was not associated with a fitness cost in a susceptible, nearly isogenic, cultivar. In competition assays in the resistant cultivar, the DNA accumulation of the TYLCV-IL clone – the parent less affected by the *Ty-1* gene in single infection – dropped below the qPCR detection level at 120 days post-infection (p.i.) and below the whitefly vector (*Bemisia tabaci*) transmissibility level at 60 days p.i. The molecular basis of the selective advantage of TYLCV-IS76 is discussed in relation to its non-canonical recombination pattern, and the RNA-dependent RNA polymerase encoded by the *Ty-1* gene.

Received 31 August 2016
Accepted 13 October 2016

INTRODUCTION

Genetic exchange via recombination has been reported frequently for both DNA and RNA viruses (Lefevre & Moriones, 2015; Wain-Hobson *et al.*, 2003; Worobey & Holmes, 1999). The variation generated by mutation and recombination constitutes the raw material on which natural selection and genetic drift act to shape populations, and is essential for virus adaptation in changing environments. Most population shifts have been reported to involve variants that differ from the preceding populations by mutations that did not necessarily involve recombination events. This is, for example, the case in population shifts

detected at the host level in patients infected with human immunodeficiency virus type 1 (HIV-1) and treated with reverse transcriptase or protease inhibitors (Condra *et al.*, 1995; Larder, 1994; Richman *et al.*, 1994); parallel mutations detected in viral variants isolated from such patients are consistent with positive selection under drug pressure. At the regional level, population shifts may be caused either by accidentally imported variants, for example the Asian 1 lineage of dengue virus (DENV) in Viet Nam (Vu *et al.*, 2010), or by variants arising locally through evolutionary processes involving positive selection or genetic drift, for example the West Nile virus WN02 genotype that displaced the NY99 genotype in the USA between 2001 and 2004 (Moudy *et al.*, 2007). Viruses of the emerging WN02 genotype were thought to have derived from the NY99 genotype by mutations. Both in DENV and West Nile virus (WNV),

A supplementary table and supplementary methods are available with the online Supplementary Material.

the selective hypothesis proposed to explain the population shift was supported by experimental results that revealed differential virus accumulation between lineages for DENV and differential efficiency of mosquito transmission between genotypes for WNV.

In some rare cases, population shifts have been reported to involve recombinant viruses (Belabess *et al.*, 2015; Monci *et al.*, 2002; Shi *et al.*, 2013), and two such cases are derived from viruses of the genus *Begomovirus* (*Geminiviridae*), i.e. TYLCV and TYLCSV of the species *Tomato yellow leaf curl virus* and *Tomato yellow leaf curl Sardinia virus*, respectively. Following introduction of the Mild strain of TYLCV (TYLCV-Mld) into Spain, two tomato-infecting begomoviruses co-circulated in this country: TYLCV-Mld and the resident Spanish strain of TYLCSV (TYLCSV-ES). One year after the first detection of the invading TYLCV-Mld in 1998, a TYLCV-Mld/TYLCSV-ES recombinant was detected: TYLCMaV (species *Tomato yellow leaf curl Malaga virus*) (Monci *et al.*, 2002). This variant displaced TYLCV-Mld viruses in bean in the Almeria province, with 76% of monitored plants (64/84) being infected only with TYLCMaV viruses. It was proposed that its extended host range, which included bean – a species that could not be infected with TYLCSV – and two non-cultivated Solanaceous species that could not be infected with TYLCV, might have provided a selective advantage to the recombinant TYLCMaV (García-Andrés *et al.*, 2007a). The second population shift associated with a recombinant was reported from Morocco with the recombinant TYLCV-IS76 (Belabess *et al.*, 2015). Here again, TYLCSV was involved as the parental virus but recombination occurred with the Israel strain of TYLCV (TYLCV-IL). TYLCV-IS76 exhibited a non-canonical recombination pattern with a 76-nt TYLCSV-derived fragment that was much shorter than that of TYLCV/TYLCSV recombinants reported previously (Davino *et al.*, 2009, 2012; García-Andrés *et al.*, 2007b; Monci *et al.*, 2002). TYLCV-IS76 was not detected between 1999 and 2003 when TYLCV-IL and TYLCSV were co-circulating in Morocco, but was found to have completely displaced its parental viruses in southern Morocco (Souss) by 2012. Interestingly, the period during which TYLCV-IS76 passed from low frequency (below detection level) to the replacement of its parental viruses in the Souss (2004–2012), coincides with the period during which susceptible tomato cultivars were replaced to a large extent with cultivars bearing the *Ty-1* resistance gene. Likewise, the partial displacement of TYLCSV by the invading TYLCV in Spain has coincided with the general deployment of *Ty-1*-resistant cultivars (García-Andrés *et al.*, 2009). In both cases, it is thought that the population shift may have been driven by the selection exerted by tomato cultivars bearing the *Ty-1* gene associated with hindrance of intra-plant viral accumulation (Michelson *et al.*, 1994). In the case of the displacement observed in Spain, the selective hypothesis has been supported by intra-plant viral DNA accumulation data showing that the *Ty-1* gene was more effective with TYLCSV than with TYLCV (García-Andrés *et al.*, 2009). In

the case of the displacement observed in Morocco, selection has been favoured over drift as the main evolutionary mechanism, since tomatoes are grown year-round in the Souss, maintaining high viral populations and preventing narrow bottlenecks.

Hence, the objective of this study was to confirm the selective hypothesis by testing, under experimental conditions, whether the fitness of TYLCV-IS76 is higher than that of representatives of its parental virus species, TYLCV-IL and TYLCSV-ES. Taking viral DNA accumulation as a proxy for fitness, we show here with real-time PCR that intra-tomato plant accumulation of TYLCV-IS76 recombinants was significantly higher than that of the parental viruses in a *Ty-1*-bearing cultivar, irrespective of the infection status (single or multiple infection) or the number of days post-infection (p.i.) at which plant samples were collected (10–120 days p.i.). Moreover, the fitness advantage in the resistant cultivar was not associated with a fitness cost in a susceptible nearly isogenic cultivar. Taken together, the results support the hypothesis that the viral population shift observed in the Souss has been driven by positive selection in the resistant tomato cultivars. The most striking feature of the TYLCV-IS76 phenotype is the drastic and deleterious effect on the accumulation of TYLCV-IL viruses, which dropped below both the quantitative PCR (qPCR) detection level and the transmissibility level, as assessed with the whitefly vector *Bemisia tabaci* (Gennadius, 1889). Using a genetically engineered virus recombinant, we show that the non-canonical recombination pattern of TYLCV-IS76 is the major determinant of its selective advantage.

RESULTS

Fitness advantage of TYLCV-IS76 in grafted and non-grafted *Ty-1*-resistant tomato plants

The population shift that resulted in the complete dominance of the recombinant TYLCV-IS76 in the Souss coincided with the increased use of *Ty-1*-resistant plants, which is thought to have positively selected for this recombinant viral strain (Belabess *et al.*, 2015). The ‘selection’ hypothesis was tested here by assessing the viral DNA accumulation of a TYLCV-IS76 clone in a *Ty-1*-resistant plant and comparing it to that of representatives of the parental species, i.e. TYLCV-IL and TYLCSV-ES clones. The *Ty-1*-resistant tomato cultivar *Pristyla* was used because it is grown in more than half of the Souss area cultivated with *Ty-1*-resistant cultivars of round-shaped tomatoes. As all the tomato plants grown by farmers in the Souss are grafted, *Pristyla* was tested not only as seedlings but also as grafted plants. *Maxifort* was used as the rootstock because it is the most commonly used rootstock in the Souss. The plants were agroinoculated with TYLCV-IS76, TYLCV-IL and TYLCSV-ES clones in various combinations.

The inoculation of 20 grafted *Pristyla* plants per viral clone was 100% successful for the three agroinfectious clones

(Table 1; experiment 1). None of the infected plants exhibited the tomato yellow leaf curl (TYLC) symptoms induced by these viral clones in susceptible tomato plants. For convenience, viral DNA accumulation of a viral clone, for example the TYLCV-IL clone, is hereafter referred to as DNA accumulation of TYLCV-IL. In competition assays, DNA accumulation of TYLCV-IS76 was significantly higher than that of both parental viruses at 30 days p.i. (Fig. 1). The most striking differences were detected between TYLCV-IS76 and TYLCV-IL, with a mean accumulation ratio [TYLCV-IS76/TYLCV-IL] of 700 in the triple infection treatment, and 500 in double infection. The mean accumulation ratio [TYLCV-IS76/TYLCV-ES] was about

10 in both triple and double infection treatments. In single infection treatments, the DNA accumulation of TYLCV-IS76 was also higher than those of TYLCV-IL and TYLCV-ES, which indicated that the fitness advantage of the TYLCV-IS76 clone revealed in competition assays is not associated with an accumulation cost in single infection. By comparing the viral DNA accumulation of each virus between single and competition infections, contrasting interactions were detected. Whereas the mean DNA accumulations of TYLCV-IS76 and TYLCV-ES were similar or higher in competition assays than in single infection, the mean DNA accumulation of TYLCV-IL in double- or triple infection with TYLCV-IS76 was more than 100 times lower

Table 1. Origin of leaf samples from which the viral DNA content was quantified by real-time PCR

Samples were collected from tomato plants of the *Ty-1*-resistant cv. Pristyla, of a susceptible nearly isogenic cultivar, and of the *Ty-1*-resistant cultivar F. Resistant cultivars were heterozygous for the *Ty-1*-resistant allele (*Ty-1/ty-1*). Plants were infected in single or mixed infection with TYLCV-IL (IL), TYLCV-ES (ES), TYLCV-IS76 (IS76) or TYLCV-IS76' (IS76') agroinfectious clones. p.i., Post-inoculation.

| Experiment | Tomato cultivar (<i>Ty-1</i> gene alleles) | Viral infection status | No. infected plants/no. inoculated plants | No. plants tested at each sampling time (days p.i.) | | | | | |
|------------|---|------------------------|---|---|----|----|----|----|-----|
| | | | | 10 | 20 | 30 | 60 | 90 | 120 |
| 1 | Pristyla (<i>Ty-1/ty-1</i>) grafted on cultivar Maxifort | IL | 20/20 | 20 | 20 | | | | |
| | | ES | 20/20 | 20 | 20 | | | | |
| | | IS76 | 20/20 | 20 | 20 | | | | |
| | | IL+ES | 20/20 | 20 | 20 | | | | |
| | | IL+IS76 | 19/20 | 19 | 19 | | | | |
| | | IS76+ES | 20/20 | 20 | 20 | | | | |
| | | IL+ES+IS76 | 34/40 | 34 | 34 | | | | |
| 2 | Pristyla (<i>Ty-1/ty-1</i>) Susceptible nearly isogenic cultivar (<i>ty-1/ty-1</i>) | IL | 33/35 | 30 | 30 | 30 | 9 | 9 | |
| | | ES | 28/35 | 28 | 28 | 28 | 9 | 9 | |
| | | IS76 | 60/60 | 30 | 30 | 30 | 10 | 10 | |
| | | IL+ES+IS76 | 39/60 | 26 | 26 | 26 | 10 | | |
| | | IL | 35/35 | 30 | 30 | 30 | 9 | 9 | |
| | | ES | 33/35 | 30 | 30 | 30 | 10 | 9 | |
| | | IS76 | 60/60 | 30 | 30 | 30 | 8 | 8 | |
| 3 | F (<i>Ty-1/ty-1</i>) | IL | 5/5 | | | 5 | | | |
| | | IS76 | 5/5 | | | 5 | | | |
| | | IL+IS76 | 5/5 | | | 5 | | | |
| 4 | Pristyla (<i>Ty-1/ty-1</i>) | IL | 15/15 | 15 | 15 | 15 | | | 8 |
| | | IL+ES | 15/15 | 15 | 15 | 15 | | | 8 |
| | | IL+IS76 | 15/15 | 15 | 15 | 15 | | | 9 |
| | | IL+ES+IS76 | 20/20 | 20 | 20 | 20 | | | 9 |
| 5 | Pristyla (<i>Ty-1/ty-1</i>) | IL | 15/15 | 13 | | 15 | | | |
| | | IS76' | 20/20 | 15 | | 20 | | | |
| | | IL+IS76 | 17/18 | 16 | | 17 | | | |
| 6 | Pristyla (<i>Ty-1/ty-1</i>) Susceptible nearly isogenic cultivar (<i>ty-1/ty-1</i>) | IL | 4/5 | 3 | | 4 | | | |
| | | IS76' | 5/5 | 5 | | 5 | | | |
| | | IL+IS76 | 5/5 | 5 | | 5 | | | |
| | | IL+IS76' | 49/51 | 15 | | 15 | | | |
| | | IL | 5/5 | 5 | | 4 | | | |
| | | IS76' | 5/5 | 5 | | 4 | | | |
| | | IL+IS76' | 20/20 | 20 | | 15 | | | |

than that in single infection. The deleterious effect on the TYLCV-IL clone was detected already at 10 days p.i., with mean DNA accumulation ratios [single/competition infections] of 4 and 8 in double and triple infection, respectively (results not shown). Interestingly, co-infection was always profitable to the DNA accumulation of TYLCSV-ES and, unlike the TYLCV-IS76 clone, TYLCSV-ES clone did not exhibit any deleterious effect on the TYLCV-IL clone.

Like for the grafted *Pristyla* plants, agroinoculation of non-grafted *Pristyla* plants was 100 % successful and did not induce any Tylc symptoms, irrespective of the agroinfectious clone (Table 1; experiment 2). The DNA accumulation patterns determined with non-grafted *Pristyla* plants were similar to those determined with grafted plants (compare Figs 1

and 2a). Thus, the DNA accumulation of TYLCV-IS76 was significantly higher than those of TYLCV-IL and TYLCSV-ES, irrespective of infection status (single or competition infections) or collection time of the samples (10, 20, 30, 60 or 90 days p.i.), which confirms the fitness advantage of TYLCV-IS76. Interestingly, whereas the differential of DNA accumulation between the three viruses tended to increase over time in the competition infection, with, for example, an increase in the mean accumulation ratio [TYLCV-IS76/TYLCV-IL] from 15 at 10 days p.i. to 3600 at 60 days p.i., the differential tended to decrease in single infections with, for example, a decrease in the mean accumulation ratio [TYLCV-IS76/TYLCSV-ES] from 1000 at 10 days p.i. to 10 at 90 days p.i. These differing accumulation dynamics between the




Fig. 1. DNA accumulations of TYLCV-IL (IL), TYLCSV-ES (ES) and TYLCV-IS76 (IS76) clones in agroinfected tomato plants of the *Ty-1*-resistant cv. *Pristyla* grafted on a tomato rootstock of the cv. *Maxifort*. The infection status of plants is indicated at the top of the figure. Viral DNA was quantified with real-time PCR from leaf samples collected at 30 days p.i. from 20 plants per treatment, except in the triple infection and the IL+IS76 double infection treatment, in which 34 and 19 plants were tested, respectively (Table 1; experiment 1). The logarithm of the Calibrated Value (logCvR) reflects viral DNA accumulation. Within the boxes, the horizontal line indicates the median value (50% quantile), the box itself delimits the 25 and 75% quantiles, and lines represent the normal range of the values; the points above and/or below correspond to outlying values. The red, blue and purple dotted lines represent the mean of logCvR values obtained with mock-inoculated plants tested with TYLCV-IL, TYLCSV-ES and TYLCV-IS76 specific primers, respectively. The red and purple dotted lines cannot be distinguished because the corresponding logCvR values are virtually the same: -6.99 and -6.97, respectively. Box plots with different letters indicate significant differences of viral DNA accumulations (Tukey's test, $P=0.05$): small letters correspond to comparisons between single-infected viruses or between different viruses in co-infected plants, whereas capital letters correspond to the comparisons of the same virus between treatments.




Fig. 2. DNA accumulations of TYLCV-IL, TYLCV-ES and TYLCV-IS76 in agroinfected non-grafted tomato plants of (a) the *Ty-1*-resistant cv. *Pristyla* (*Ty-1/ty-1*) and (b) a susceptible nearly isogenic cultivar (*ty-1/ty-1*). The infection status of plants, single or triple infection, is indicated at the top of the figure. Viral DNA was quantified with real-time PCR from leaf samples collected between 10 and 90 days p.i. from at least 26 plants for the 10, 20 and 30 days p.i. samplings, and from at least 8 plants for the 60 and 90 days p.i. samplings (Table 1; experiment 2). Representation of viral DNA accumulations, box plots, positive thresholds and statistics as in Fig. 1. Small letters correspond to the differences between the three viruses among samples collected at the same time from plants of the same cultivar, and with the same infection status, whereas capital letters correspond to the comparisons of the same virus between single- and triple-infected plants of the same cultivar and sampled at the same time.

single and competition infections were due mainly to the DNA accumulation of TYLCV-IL in triple-infected plants, which was drastically lower than that in single-infected plants, irrespective of sampling time. The deleterious effect was strong enough to be detectable at an early stage of infection, with a TYLCV-IL DNA accumulation ratio [single/competition infection] of 10 at 10 days p.i.; the deleterious effect apparently did not decrease over time according to the 60 days p.i. ratio of mean accumulations, which was 1500. Unlike DNA accumulation of TYLCV-IL, DNA accumulations of TYLCV-IS76 and TYLCV-ES were not much affected by co-infection, and their accumulations were not significantly different between single- and triple-infected plants at the latest common collection date, i.e. 60 days p.i.

Taken together, the DNA accumulation patterns of TYLCV-IS76 and parental viruses in *Pristyla* plants (Figs 1 and 2a) were consistent with the hypothesis of a critical role played by *Ty-1*-resistant plants in the emergence, and eventually the complete dominance, of TYLCV-IS76 viruses in the Souss.

The fitness advantage of TYLCV-IS76 in *Ty-1*-resistant plants is not associated with a fitness cost in susceptible plants

To further confirm the selective hypothesis of displacement of the parental viruses by TYLCV-IS76 recombinant viruses in the Souss agroecosystem, we tested if the fitness advantage of TYLCV-IS76 in *Ty-1*-resistant plants could be associated with any cost with respect to its DNA accumulation in susceptible non-*Ty-1*-bearing plants. Hence, the DNA accumulations of the three viruses were tested in a susceptible cultivar that is nearly isogenic with *Pristyla*. For this and the following tests, non-grafted tomato plants were used as a more straightforward model that did not require the assistance of a commercial nursery. This simplification was possible because, as reported above, the fitness advantage of TYLCV-IS76 seen in grafted *Pristyla* plants was also found in non-grafted *Pristyla* plants (compare Figs 1 and 2a). Thus, the resistant and susceptible non-grafted tomato plants were agroinoculated in the same experiment with the TYLCV-IS76, TYLCV-IL and

TYLCSV-ES clones, and leaf samples were collected between 10 and 90 days p.i. (Table 1; experiment 2).

The Tylc symptoms induced by TYLCV-IS76 in plants of the susceptible cultivar were similar to those induced by TYLCV-IL and TYLCSV-ES. The viral DNA contents of the susceptible plants were significantly higher than those of the resistant plants for the three viruses (compare Fig. 2a and Fig. 2b, $P < 2.2 \times 10^{-16}$, for single- and triple-infected plants). The DNA accumulation of TYLCV-IS76 in the susceptible cultivar was either higher than those of both parental viruses or similar to that of the parental virus exhibiting the highest DNA accumulation (Fig. 2b). Hence, the fitness advantage of TYLCV-IS76 in *Pristyla* was not associated with an apparent cost with respect to its accumulation in the susceptible cultivar. In spite of its high accumulation in the susceptible plants, TYLCV-IS76 DNA accumulation was relatively less affected than those of parental viruses in the *Ty-1*-resistant cultivar. Indeed, the mean DNA accumulation ratios between susceptible plants and resistant plants were always lower for TYLCV-IS76 than for parental viruses, irrespective of infection status, and date of sampling, except in the single infection treatment at 30 days p.i. where the ratios were similar between TYLCV-IL and TYLCV-IS76 clones, around 29. Interestingly, the mean DNA accumulation ratios [TYLCV-IS76/parental virus (TYLCV-IL or TYLCSV-ES)] were highest in samples of the earliest collection dates (10 or 20 days p.i.) in both the competition and single infection assays in the resistant cultivar, which suggests that TYLCV-IS76 has an advantage at the onset of infection. It is noteworthy that competition is deleterious to the accumulation of TYLCV-IL DNA in susceptible plants although not as drastically as in resistant plants. Indeed, the ratio of TYLCV-IL DNA mean accumulations between single- and triple-infected plants at 10, 30 and 60 days p.i. was 10, 10 and 80 in susceptible plants, and 11, 42 and 1500 in resistant plants, respectively.

The fitness advantage of TYLCV-IS76 is related to the *Ty-1* resistance gene

To test if the fitness advantage of TYLCV-IS76 viruses in *Pristyla* was due mainly to the *Ty-1* gene rather than to the genetic background of the cultivar *Pristyla* or its specific interaction with the *Ty-1* gene, we used the cultivar 'F', which has the same *Ty-1* allele as *Pristyla* but in a different genetic background. As the major feature of the fitness advantage of TYLCV-IS76 was its deleterious effect on DNA accumulation of TYLCV-IL, the test was limited to three treatments: single infection with the TYLCV-IL clone, single infection with the TYLCV-IS76 clone, and a competition test with both these clones; five F plants were infected for each treatment and sampled at 30 days p.i. (Table 1; experiment 3). DNA accumulation of TYLCV-IS76 was significantly higher than that of TYLCV-IL in the single and competition infections (Wilcoxon test, $P = 0.007$) (Fig. 3), and the mean accumulation ratios (TYLCV-IS76/TYLCV-IL) were of the same order of

magnitude as those detected with *Pristyla*, i.e. 200 in co-infected plants and 6 in single-infected plants. These results support the view that the selective advantage of TYLCV-IS76 over TYLCV-IL was due primarily to the *Ty-1* gene that both cultivars have in common, and only marginally dependent (if at all) on their different genetic background.

From 60 days p.i., *Pristyla* tomato plants tend to be dead ends for TYLCV-IL when co-infected with TYLCV-IS76

The deleterious effect of TYLCV-IS76 viruses on TYLCV-IL viruses in *Pristyla* plants was observed until 60 days p.i. (Fig. 2a). To test if it is maintained beyond this date, and might result in the elimination of TYLCV-IL viruses, viral DNA accumulations were further monitored up to 120




Fig. 3. DNA accumulations of TYLCV-IL (IL) and TYLCV-IS76 (IS76) in agroinfected non-grafted tomato plants of the *Ty-1*-resistant cv. F (*Ty-1/ty-1*). The infection status of plants, single or double infection, is indicated at the top of the figure. Viral DNA was quantified with real-time PCR from leaf samples collected at 30 days p.i. from five plants per treatment (Table 1; experiment 3). Representation of DNA accumulations, box plots and positive thresholds as in Fig. 1.

days p.i. (Table 1; experiment 4). Fifteen *Pristyla* plants double-infected with the TYLCV-IL and TYLCV-IS76 clones, and 20 plants triple-infected with the TYLCV-IL, TYLCV-IS76 and TYLCSV-ES clones were sampled at 10, 20 and 30 days p.i., and nine of them were sampled randomly at 120 days p.i. Fifteen plants infected with the TYLCV-IL clone alone and 15 plants co-infected with the TYLCV-IL and TYLCSV-ES clones were tested in parallel as negative controls in which TYLCV-IL viruses were not subjected to the deleterious effect of TYLCV-IS76 viruses. As in the previous experiments (Figs 1–3), DNA accumulation of TYLCV-IL was drastically affected by the presence of TYLCV-IS76 in both the double- and triple-infected plants (Fig. 4). Thus, at 30 days p.i., DNA accumulation of TYLCV-IL in these plants was more than 200 times lower than that of TYLCV-IL in plants without TYLCV-IS76. Interestingly, the amount of TYLCV-IL DNA dropped below the detection level at 120 days p.i. in plants co-infected with TYLCV-IS76 in both competition treatments;

it remained detectable in only one of the nine triple-infected plants tested. In the plants infected only with TYLCV-IL viruses, the mean DNA accumulation at 120 days p.i. was similar to that at 30 days p.i. According to the 120 days p.i. samplings, it seems that TYLCSV-ES has also a negative impact on the DNA accumulation of TYLCV-IL, as the mean DNA accumulation of TYLCV-IL in competition with TYLCSV-ES was significantly lower than that of TYLCV-IL in single infection.

It was expected that the negative impact of TYLCV-IS76 viruses on TYLCV-IL viruses may compromise the transmission of TYLCV-IL from co-infected plants by the whitefly vector *B. tabaci*. A preliminary transmission test was conducted with two *Pristyla* plants in which the DNA accumulation of TYLCV-IS76 were 2500 and 4000 times higher than that of TYLCV-IL, respectively, at 56 days p.i. TYLCV-IS76 was the only virus detected among infected test plants (6/15 plants of the susceptible cultivar and 3/15 plants of the resistant cultivar).




Fig. 4. DNA accumulations of TYLCV-IL in non-grafted tomato plants of the *Ty-1*-resistant cv. *Pristyla*, either single-infected or co-infected with TYLCV-IS76 (IS76), TYLCSV-ES (ES) or both. The infection status of plants is indicated at the top of the figure. Viral DNA was quantified with real-time PCR from leaf samples collected between 10 and 120 days p.i. from at least 15 plants for the 10, 20 and 30 days p.i. samplings, and from at least 8 plants for the 120 days p.i. samplings (Table 1; experiment 4). Representation of viral DNA accumulations, box plots, positive thresholds and statistics as in Fig. 1. Different letters indicate significant differences of TYLCV-IL DNA accumulations between plant samples of the four treatments collected at the same time.

Recombination is the determinant of the antagonistic impact of TYLCV-IS76 on TYLCV-IL

It was inferred from Bayesian analysis that the recombination event leading to this TYLCV-IS76 viruses occurred most probably in the late 1990s (Belabess *et al.*, 2015). Therefore, as the genuine parents of TYLCV-IS76 viruses were not available, the clones used as representatives of the parental species in the experimental studies were selected among those that exhibited the highest nucleotide identity with them. Thus, the TYLCV-IL-derived fragment genome of the TYLCV-IS76 clone differed from that of the selected TYLCV-IL clone by 27 mutations. Conversely, the TYLCV-ES-derived fragment genome of the TYLCV-IS76 clone differed from that of the selected TYLCV-ES clone by two mutations. Hence, to determine if recombination alone may have provided a fitness advantage to the TYLCV-IS76 recombinant over its genuine and unknown parental TYLCV-IL, we used the agroinfectious clone TYLCV-IL to engineer TYLCV-IS76' - a recombinant with the same recombination profile as TYLCV-IS76 but which is 100% identical to the genome of its parental TYLCV-IL in its TYLCV-IL-derived region and 100% identical to the genome of the clone TYLCV-IS76 in its TYLCV-ES derived region.

The infectivity of the engineered TYLCV-IS76' was similar to that of its parental clone TYLCV-IL in both cultivars: TYLCV-IS76' and TYLCV-IL infected 100% of inoculated *Pristyla* plants (20 and 15 inoculated plants, respectively), and 100% of nearly isogenic susceptible plants (5 for each clone) (Table 1; experiments 5 and 6). TYLCV-IS76' produced the typical Tylc symptoms of TYLCV-IL in the susceptible cultivar. Moreover, in single-infected resistant plants, the DNA accumulation of TYLCV-IS76' was similar to that of TYLCV-IL at 30 days p.i. (Fig. 5), which, altogether indicates that recombination at position 76 had no negative impact on infectivity or viral DNA accumulation of the engineered recombinant in comparison to its TYLCV-IL progenitor. Consistently, the sequence of four full-length TYLCV-IS76' genomes cloned from a 30 days p.i. plant co-infected with the TYLCV-IL clone were 100% identical to that of the agroinoculated clone. In competition tests, the DNA accumulation of TYLCV-IS76' was significantly higher than that of its progenitor TYLCV-IL in both cultivars at 30 days p.i. Taken together, and, similarly to the wild-type TYLCV-IS76 (Figs 1–3), the fitness advantage of the TYLCV-IS76' revealed in competition assays with TYLCV-IL is not associated with a cost with respect to its accumulation in single infection. As a control, *Pristyla* plants were co-infected in parallel with the wild-type TYLCV-IS76 and TYLCV-IL (Fig. 5). Interestingly, the DNA accumulation of TYLCV-IL was similarly low in both competition tests, indicating that the 27 discriminating mutations between the engineered and the wild-type recombinants had no significant impact on the deleterious effect induced by recombination. However, the slight but significant difference in virus accumulation between the natural Moroccan recombinant and the recombinant

engineered with a TYLCV-IL clone from Réunion is determined obviously by mutations (Fig. 5).

DISCUSSION

The TYLCV-IS76 recombinants that have emerged in the Souss region of southern Morocco have virtually replaced




Fig. 5. DNA accumulations of TYLCV-IL (IL), TYLCV-IS76 (IS76) and TYLCV-IS76' (IS76') in agroinfected non-grafted tomato plants of the *Ty-1*-resistant cv. *Pristyla* (*Ty-1/ty-1*) and a susceptible nearly isogenic cultivar (*ty-1/ty-1*). The infection status of the plants is indicated at the top of the figure. Viral DNA was quantified with real-time PCR from leaf samples collected at 10 and 30 days p.i. from 15–20 plants per treatment (Table 1; experiments 5 and 6). As there was no statistical difference between the results of experiments 5 and 6 (Table 1), the data obtained for the samples collected at 10 and 30 days p.i. of these two experiments were pooled for statistical analysis. Representation of viral DNA accumulations, box plots, positive thresholds and statistics as in Fig. 1. The positive threshold for TYLCV-IS76' and TYLCV-IS76 is the same because the same primer pair was used for both viruses. Small letters correspond to the differences between different viruses among samples collected at the same time from plants of the same cultivar and with the same infection status, capital letters correspond to the comparisons of the same virus between single- and double-infected plants of the same cultivar and sampled at the same time, and bold capital letters correspond to the comparisons of IS76 and IS76' virus in the resistant cultivar and sampled at the same time.

their parental viruses since the deployment of *Ty-1*-cultivars in the 2000s. The experimental fitness comparisons of the recombinant and parental type viruses presented here provide compelling results to support the selective advantage of TYLCV-IS76 over parental type viruses and, most importantly, its selection by *Ty-1*-cultivars without any detectable fitness cost in non-*Ty-1*-bearing cultivars.

Measuring nucleic acid accumulation has already been used to show the fitness advantage of a cucumber mosaic virus recombinant produced under greenhouse conditions (Fernández-Cuartero *et al.*, 1994). However, although natural emerging recombinants have been previously reported (Davino *et al.*, 2009; García-Andrés *et al.*, 2006; Monci *et al.*, 2002; Shi *et al.*, 2013; Zhou *et al.*, 1997), a mechanistic basis for such population shifts has only been proposed with TYLCMaIV, the recombinant which partially displaced its parental viruses in bean. It was reported to have a slightly higher infectivity in *Ty-1*-resistant tomato plants, and an extended host range in comparison with those of representatives of its parental species (Monci *et al.*, 2002). However, its transmission efficiency from single-infected plants was not higher than that of its parents, and its viral DNA accumulation assessed from dot blot hybridizations was only 14 % of TYLCV - the parent exhibiting the highest accumulation. Similar results were obtained with two TYLCV-IL/TYLCSV-Sar recombinant viruses from Italy exhibiting the same recombination profiles as TYLCaXV – the other TYLCV/TYLCSV recombinant from Spain (García-Andrés *et al.*, 2006) – and TYLCMaIV; their transmission efficiency was not higher than that of the parental viruses, and their replication in agroinfiltrated *Nicotiana benthamiana* leaves was estimated to be about 10 times lower than that of a representative clone of the parental TYLCSV species (Davino *et al.*, 2009). The low accumulation of these canonical recombinants may explain why plants infected only with such recombinants were not detected in Italy, and why the replacement of parental viruses by recombinants has not been reported from Spain.

According to a host range study on tomato, common bean (cv. Contender) and *Solanum nigrum*, the TYLCV-IS76 clone did not exhibit any extended host range compared to that of the parental clones (data not shown).

Contrasted efficiency of the *Ty-1* gene to control TYLCV-IS76 and its parental viruses

The viral DNA contents of the susceptible plants were significantly higher than those of the resistant plants for the three viruses (compare Fig. 2a and Fig. 2b). This result is consistent with the reported negative impact of the *Ty-1* resistance gene on viral DNA accumulation previously monitored by southern or dot-blot hybridization (Barbieri *et al.*, 2010; García-Andrés *et al.*, 2009; Michelson *et al.*, 1994) and monitored here with qPCR for the first time. More specifically, the *Ty-1* gene was more effective to restrain the TYLCSV-ES clone than the TYLCV-IL clone which is consistent with squash-blot results obtained

previously with representatives of TYLCSV-ES and TYLCV-Mld (García-Andrés *et al.*, 2009). TYLCV-IS76 was the less affected virus in the resistant cultivar, particularly in competition tests. Moreover, it is noteworthy that the deleterious effect of TYLCV-IS76 on TYLCV-IL DNA accumulation observed is much higher in *Pristiella* than in susceptible plants (Fig. 2), which is consistent with the supposed triggering effect of *Ty-1*-resistant plants on the emergence of TYLCV-IS76 recombinants in Morocco.

Recombinant and parental viruses exhibit contrasting viral DNA accumulation and competitiveness

Comparison of virus accumulation between single- and multiple-infected plants revealed that the three viral clones were affected differently by competition. Competition was always deleterious for DNA accumulation of TYLCV-IL (Figs 1 and 2) except in the double infection with TYLCSV-ES at 30 days p.i. (Fig. 1). On the contrary, competition was always beneficial or neutral to DNA accumulation of TYLCSV-ES, except in triple-infected susceptible plants at 10 days p.i. (Fig. 2). These results are consistent with the relatively higher frequency of TYLCSV than TYLCV-Mld amplicons generated and cloned from 400 days p.i. samples of tomato plants of susceptible (García-Andrés *et al.*, 2007b) and *Ty-1*-resistant (García-Andrés *et al.*, 2009) cultivars co-infected with these viruses.

Competition is mostly beneficial or neutral to DNA accumulation of TYLCV-IS76 except at the earliest sampling times, i.e. 10 and 20 days p.i. in the susceptible cultivar and 10 days p.i. in the resistant cultivar (Fig. 2). Interestingly, a positive cooperative interaction was detected with the TYLCSV-ES clone at 30 days p.i. in the *Ty-1*-cultivar (Figs 1 and 2a). Similar synergies in DNA accumulations have been detected with two begomoviruses infecting tomato in India (Chakraborty *et al.*, 2008), as well as two begomoviruses of cassava (Fondong *et al.*, 2000).

The negative impact of the TYLCSV-ES clone on the DNA accumulation of the TYLCV-IL clone and its positive cooperative interaction with the TYLCV-IS76 clone, are consistent with the higher number of TYLCSV (6) than TYLCV-IL infected plants (1) within the 301 Tylc virus-positive tomato plants sampled in the Sous after the displacement of parental viruses by the recombinant virus (Belabess *et al.*, 2015).

Methylation may explain the low fitness of TYLCV-IL in competition tests

The 76-nt TYLCSV-ES-derived region of TYLCV-IS76 was identified as the molecular determinant of the dramatic deleterious effect on the DNA accumulation of TYLCV-IL clone in the *Ty-1*-resistant cultivar. Although no particular function has been associated to this region in geminiviruses, it has been shown to be one of the favoured regions for siRNA (small interfering RNA) targeting and methylation

in the case of the A component of three bipartite begomoviruses, mungbean yellow mosaic India virus (MYMIV) (Yadav & Chattopadhyay, 2011), tomato leaf curl New Delhi virus (ToLCNDV) (Sahu *et al.*, 2014), and pepper golden mosaic virus (PepGMV) (Rodríguez-Negrete *et al.*, 2009). Intriguingly, this was not the case of the intergenic region of the monopartite begomoviruses, tomato yellow leaf curl China virus (Yang *et al.*, 2011) and TYLCV (Butterbach *et al.*, 2014). However, as the DNA accumulation of TYLCV-IL was affected in co-infection experiments, with both TYLCSV-ES and TYLCV-IS76, it is possible that co-infection mimics the infection with a bipartite begomovirus. Thus, the 76-nt discriminating nucleotides of TYLCV-IL genome may be targeted by silencing mechanisms triggered by the co-infecting viruses. As the level of silencing may depend on the concentration of the co-infecting virus, it is supposed to be higher with TYLCV-IS76 whose DNA accumulation is 10 times higher than that of TYLCSV-ES. The silencing hypothesis is consistent with the results of Butterbach *et al.* (2014) showing that the *Ty-1* resistance against TYLCV is associated with increasing cytosine methylation of the viral genome suggestive of enhanced transcriptional gene silencing (TGS). The involvement of TGS in the deleterious effect of TYLCV-IS76 on TYLCV-IL DNA accumulation should be tested experimentally.

Selection-driven displacement scenario of Tylc-associated viruses leading to the dominance of TYLCV-IS76 viruses in the Sous

According to previous reports and the results presented here on the fitness advantage of TYLCV-IS76, the following scenario is proposed to account for the replacement of parental viruses by the recombinant TYLCV-IS76 viruses in southern Morocco (Sous). At the beginning of the 2000s, when the susceptible cultivars started to be replaced with *Ty-1*-cultivars in Morocco, the population of viruses associated with Tylc symptoms was composed mainly of TYLCV-IL, TYLCV-Mld and TYLCSV viruses, and some canonical TYLCV-IL/TYLCSV recombinants, but TYLCV-IS76 viruses had not been detected (Belabess *et al.*, 2015). TYLCSV viruses are thought to have been displaced by TYLCV-Mld viruses, because according to field surveys conducted in Spain and experimental studies, TYLCV-Mld has a better ecological performance than TYLCSV (Sánchez-Campos *et al.*, 1999), and was shown to accumulate to a greater extent than TYLCSV in *Ty-1*-cultivars (García-Andrés *et al.*, 2009). TYLCV-Mld viruses are thought to have been displaced by TYLCV-IL viruses because the non-recombinant viruses detected in the surveys conducted in northern Morocco between 2008 and 2014, were mostly TYLCV-IL viruses (Belabess *et al.*, 2015), which is consistent with surveys conducted in Réunion, where the introduction of TYLCV-IL viruses has displaced the resident TYLCV-Mld viruses (Pérefarres *et al.*, 2014). According to Bayesian inferences, the recombination event that led to TYLCV-IS76 viruses most probably occurred at the end of the 1990s, which suggests that

they were already present at the beginning of the 2000s but probably at low prevalence, at least in tomato (Belabess *et al.*, 2015). Then, in accordance with the results of the present study, where a TYLCV-IS76 clone was shown to have a selective advantage over parental clones and particularly a TYLCV-IL clone, TYLCV-IL viruses have been displaced by TYLCV-IS76 viruses. The complete displacement of TYLCV-IL suggests that the selective advantage of TYLCV-IS76 was highly efficient, which is fully consistent with the drastic deleterious effect of both TYLCV-IS76-type recombinants (IS76 and IS76') on the DNA accumulation of TYLCV-IL in co-infected *Ty-1*-plants (Figs 4 and 5). Moreover, as suggested by a preliminary transmission test, the transmission of TYLCV-IL from co-infected *Ty-1*-plants seems to be compromised, which is fully consistent with the previously reported correlation between viral DNA accumulation and transmission efficiency of Tylc-associated viruses (Lapidot *et al.*, 2001).

Hence, the co-occurrence of the replacement of susceptible cultivars by resistant ones and the complete displacement of Tylc-associated viruses by TYLCV-IS76 viruses may be more than a coincidence. Indeed, as the selective advantage of TYLCV-IS76 was far higher in *Ty-1* than in susceptible cultivars in both single and competition tests, our results strongly support the hypothesis that the deployment of *Ty-1*-cultivars has played a role in the observed viral population shift.

Although the results presented here provide compelling support for the selective hypothesis to explain the entire displacement of the parental viruses by the TYLCV-IS76 recombinants, no new light is shed on the origin of the non-canonical TYLCV-IS76 recombinant. The reasons why it was detected in Morocco but not in other countries where TYLCV/TYLCSV recombinants are frequently reported (Spain and Italy) are presently not clear. As a first step, it will be useful to test if TYLCV-IS76 viruses can be generated in tomato plants co-infected with representatives of its two parental species, and, if that is the case, would the resistant plants bearing the *Ty-1* gene be more conducive to an increase of their intra-plant frequency in comparison to susceptible plants.

METHODS

Plant material. Nearly isogenic susceptible and *Ty-1*-resistant cultivars of tomato were used to test the effect of the *Ty-1* gene on virus accumulation. The resistance conferred by this gene was recently showed to act through TGS (Butterbach *et al.*, 2014). The resistant cultivar 'Pristyla' carrying the *Ty-1* resistance allele in a heterozygote state (*Ty-1/ty-1*) is a hybrid cultivar obtained by crossing a *Ty-1*-bearing line Ar (*Ty-1/Ty-1*) with a susceptible line B; the line Ar was derived from the susceptible line A following *Ty-1* introgression. The nearly isogenic susceptible cultivar of Pristyla is the hybrid A x B. The effect of the *Ty-1* gene was tested also with the resistant cultivar 'F', which has the same *Ty-1* gene with Pristyla but introgressed in a different genetic background; the *Ty-1* allele of F was also in a heterozygote state. The term 'resistant' was chosen to designate *Ty-1*-bearing cultivars because the *Ty-1* gene induces resistance to TYLCV DNA accumulation (Pérez de Castro *et al.*, 2005; Michelson *et al.*, 1994). Seven-day-old seedlings were transplanted into

individual pots for the tests. Grafted and non-grafted *Pristiyla* plants were tested. Grafting was performed by Nimaplants nursery (Nîmes, France) 20 and 22 days after sowing *Pristiyla* and the rootstock 'Maxi-fort', respectively.

All plants were grown in containment growth chambers under 14 h light at 26±2 °C, and 10 h dark at 24±2 °C, and were watered with 15:10:30 NPK fertilizer + oligoelements.

Agroinfectious clones. Three agroinfectious clones were used: a clone of the recombinant TYLCV-IS76, and a clone of each of the two parental viruses, TYLCV-IL and TYLCSV-ES. The agroinfectious clone TYLCV-IS76[MA:SouG8:10] (GenBank accession number LN812978) has been described previously (Belabess *et al.*, 2015). Representatives of the parental species were selected among those which exhibited the highest nucleotide identity with the TYLCV- and TYLCSV-derived sequences of the recombinant TYLCV-IS76. The clone TYLCV-IL[RE:STG4:04] (GenBank accession number AM409201) (Belabess *et al.*, 2015) exhibits 99% nucleotide identity with the TYLCV-derived fragment of the TYLCV-IS76 clone. An agroinfectious clone was constructed previously as follows. A 0.9-mer genome obtained by digestion with *NcoI* and *EcoRI* was ligated into the corresponding restriction sites of the vector pCAMBIA0380. The full-length genome was excised from the plasmid pGEMT with *NcoI* and ligated into the *NcoI* restriction site of the recombinant pCAMBIA0380. The construction was introduced into bacteria of the C58 MP90 strain of *Agrobacterium tumefaciens* via electroporation. The clone TYLCSV-ES[MA:Aga5a:12] (GenBank accession number LN846598) (Belabess *et al.*, 2015) differs at only two nucleotide positions (32 and 44) from the TYLCSV-derived region of the TYLCV-IS76 clone; an agroinfectious clone has been constructed previously (Belabess *et al.*, 2015).

Construction of recombinant TYLCV-IS76' by site-directed mutagenesis. The recombinant TYLCV-IS76' was engineered starting from TYLCV-IL[RE:STG4:04] with the QuikChange Lightning Site-Directed Mutagenesis Kit (Agilent Technologies) and primers (Table S1, available in the online Supplementary Material). An agroinfectious clone of TYLCV-IS76' was prepared according to the partial tandem method described above.

Agroinoculation and randomized experimental design. Tomato plants were agroinoculated 14 days after sowing or grafting with various combinations of agroinfectious clones as described in Table 1. The transformed *A. tumefaciens* clones were grown at 28 °C in liquid LB medium containing kanamycin (50 mg ml⁻¹) and gentamycin (20 mg ml⁻¹). After about 26 h, when suspensions reached optical densities (OD_{600 nm}) of about 3–5, the agrobacteria cultures containing the infectious TYLCV-IL, TYLCSV-ES and TYLCV-IS76 genomes were adjusted to identical ODs with LB medium. The equally concentrated cultures were centrifuged for 20 min at 1000 g, and each pellet was resuspended in water (the same volume as the centrifuged volume) containing 150 µM acetosyringone and 10 mM MgCl₂. For mixed virus infections, the same procedure was used except that the same volumes of equally concentrated agrobacteria cultures containing the infectious viral genomes were mixed before centrifugation. The resuspension volume of the mixed inocula was such that each virus was inoculated at the same agrobacterial concentration in single and mixed infections. Non-grafted plants (Table 1; experiments 2–6) were agroinfiltrated via transepidermal delivery of the agrobacterial suspension using a needle-less syringe applied to the underside of cotyledons. The young leaves of grafted plants (Table 1; experiment 1) were scratched at the injection spot before agroinfiltration. Some plants used as negative controls were agroinfiltrated with bacteria of the C58 MP90 strain of *A. tumefaciens* containing an empty pCAMBIA2300 plasmid. Plants were arranged in a complete randomized block design.

Total DNA extraction. Leaf samples collected from each plant at each collection date were taken from the youngest leaf for which five leaflets were visible, and consisted of five 4-mm diameter leaf discs, one per leaflet. Total DNA from each sample was extracted according to the protocol of Dellaporta *et al.* (1983) with previously reported modifications (Urbino *et al.*, 2013), and stored at -20 °C until use.

Real-time PCR quantification of each virus. The content of intraplant viral DNA was determined with real-time PCR (qPCR) and was used as a proxy for fitness, as previously applied to TYLCV (Pérefarres *et al.*, 2014; Urbino *et al.*, 2013; Vuillaume *et al.*, 2011) and other viruses (Carrasco *et al.*, 2007; Gómez *et al.*, 2009; Hillung *et al.*, 2015; Tromas *et al.*, 2014). Primer pairs were designed on both sides of the origin of replication (OR) for the specific detection of each of the parental clones (TYLCV-IL and TYLCSV-ES), and on both sides of locus 76 for detection of the TYLCV-IS76 and TYLCV-IS76' clones (Table S1). The specificity of each primer pair was tested with viral DNA of non-targeted viral clones. Moreover, the primer pair targeting recombinants did not produce any positive detection with DNA extracts from plants double-infected with TYLCV-IL and TYLCSV clones, or with a mix of plasmid DNA extracts of the parental viral clones, indicating that no recombinants were generated *in vitro*.

The viral DNA content of each agroinfected plant was quantified in duplicate using the LightCycler 480 SYBR Green I Master qPCR mix (Roche) as described in the online Supplementary Material. All PCR fluorescence data were analysed as described in the Supplementary Material.

Transmission tests. Transmission tests were carried out with Q1 type *B. tabaci* whiteflies of the putative species Mediterranean (Med) (Angers, France), which is the most common species in Morocco (Tahiri *et al.*, 2006; 2013). Approximately 300 adults newly emerged were given a 2-day acquisition access period (AAP) on two *Ty-1*-resistant tomato plants (cv. *Pristiyla*), 56 days after their co-agroinfection with the TYLCV-IL and TYLCV-IS76 clones (Table 1; experiment 6). At the end of the AAP, 120 whiteflies were shifted to 16-day-old *Pristiyla* plants (15 plants) for a 5-day inoculation access period (IAP), and similarly 120 whiteflies were shifted to 15 plants of the susceptible cultivar. The transmission success of each virus was assessed with Multiplex PCR tests (Belabess *et al.*, 2015), which can distinguish between TYLCV-IL and TYLCV-IS76 viruses.

Statistical analysis. All statistical analyses were performed using R Studio software, version 3.0.3. (R_Development_Core_Team, 2010). Viral DNA accumulations were compared between or within plants using the log transformation of CVr data. As there was no statistical difference between the results of experiments 5 and 6 (Table 1), the data obtained for the samples collected at 10 and 30 days p.i. of these two experiments were pooled for statistical analysis. ANOVA tests were used to analyse virus accumulation data for statistical significance. A non-parametric test, Wilcoxon test, was performed with data from experiment 3 because of the low numbers of plants.

ACKNOWLEDGEMENTS

We thank Sophie Le Blaye and Jean-Luc Macia (BGPI, INRA Montpellier) for technical assistance. We also thank Stéphane Blanc (BGPI, INRA Montpellier) for useful comments and suggestions on the manuscript. Z. B., as a PhD student of the Faculté des Sciences de Meknès, received a grant from Centre National de la Recherche Scientifique et Technique (CNIRST) of Morocco. This research was supported by Gautier Semences (France) and CIRAD (<http://www.cirad.fr/>).

REFERENCES

- Barbieri, M., Acciarri, N., Sabatini, E., Sardo, L., Accotto, G. & Pecchioni, N. (2010). Introgression of resistance to two Mediterranean virus species causing tomato yellow leaf curl into a valuable traditional tomato variety. *J Plant Pathol* **92**, 485–493.
- Belabess, Z., Dallot, S., El-Montaser, S., Granier, M., Majde, M., Tahiri, A., Blenzar, A., Urbino, C. & Peterschmitt, M. (2015). Monitoring the dynamics of emergence of a non-canonical recombinant of Tomato yellow leaf curl virus and displacement of its parental viruses in tomato. *Virology* **486**, 291–306.
- Butterbach, P., Verlaan, M. G., Dullemans, A., Lohuis, D., Visser, R. G., Bai, Y. & Kormelink, R. (2014). Tomato yellow leaf curl virus resistance by *Ty-1* involves increased cytosine methylation of viral genomes and is compromised by cucumber mosaic virus infection. *Proc Natl Acad Sci U S A* **111**, 12942–12947.
- Carrasco, P., de la Iglesia, F. & Elena, S. F. (2007). Distribution of fitness and virulence effects caused by single-nucleotide substitutions in Tobacco Etch virus. *J Virol* **81**, 12979–12984.
- Chakraborty, S., Vanitharani, R., Chattopadhyay, B. & Fauquet, C. M. (2008). Supervirulent pseudorecombination and asymmetric synergism between genomic components of two distinct species of begomovirus associated with severe tomato leaf curl disease in India. *J Gen Virol* **89**, 818–828.
- Condra, J. H., Schleif, W. A., Blahy, O. M., Gabryelski, L. J., Graham, D. J., Quintero, J. C., Rhodes, A., Robbins, H. L., Roth, E. & other authors (1995). *In vivo* emergence of HIV-1 variants resistant to multiple protease inhibitors. *Nature* **374**, 569–571.
- Davino, S., Napoli, C., Dellacroce, C., Miozzi, L., Noris, E., Davino, M. & Accotto, G. P. (2009). Two new natural begomovirus recombinants associated with the tomato yellow leaf curl disease co-exist with parental viruses in tomato epidemics in Italy. *Virus Res* **143**, 15–23.
- Davino, S., Miozzi, L., Panno, S., Rubio, L., Davino, M. & Accotto, G. P. (2012). Recombination profiles between tomato yellow leaf curl virus and tomato yellow leaf curl Sardinia virus in laboratory and field conditions: evolutionary and taxonomic implications. *J Gen Virol* **93**, 2712–2717.
- Dellaporta, S. L., Wood, J. & Hicks, J. B. (1983). A plant DNA miniprep-paration: Version II. *Plant Mol Biol Report* **1**, 19–21.
- Fernández-Cuartero, B., Burguán, J., Aranda, M. A., Salánki, K., Moriones, E. & García-Arenal, F. (1994). Increase in the relative fitness of a plant virus RNA associated with its recombinant nature. *Virology* **203**, 373–377.
- Fondong, V. N., Pita, J. S., Rey, M. E., de Kochko, A., Beachy, R. N. & Fauquet, C. M. (2000). Evidence of synergism between African cassava mosaic virus and a new double-recombinant geminivirus infecting cassava in Cameroon. *J Gen Virol* **81**, 287–297.
- García-Andrés, S., Tomás, D. M., Navas-Castillo, J. & Moriones, E. (2009). Resistance-driven selection of begomoviruses associated with the tomato yellow leaf curl disease. *Virus Res* **146**, 66–72.
- García-Andrés, S., Monci, F., Navas-Castillo, J. & Moriones, E. (2006). Begomovirus genetic diversity in the native plant reservoir *Solanum nigrum*: evidence for the presence of a new virus species of recombinant nature. *Virology* **350**, 433–442.
- García-Andrés, S., Accotto, G. P., Navas-Castillo, J. & Moriones, E. (2007a). Founder effect, plant host, and recombination shape the emergent population of begomoviruses that cause the tomato yellow leaf curl disease in the Mediterranean basin. *Virology* **359**, 302–312.
- García-Andrés, S., Tomás, D. M., Sánchez-Campos, S., Navas-Castillo, J. & Moriones, E. (2007b). Frequent occurrence of recombinants in mixed infections of tomato yellow leaf curl disease-associated begomoviruses. *Virology* **365**, 210–219.
- Gómez, P., Sempere, R. N., Elena, S. F. & Aranda, M. A. (2009). Mixed infections of Pepino mosaic virus strains modulate the evolutionary dynamics of this emergent virus. *J Virol* **83**, 12378–12387.
- Hillung, J., Cuevas, J. M. & Elena, S. F. (2015). Evaluating the within-host fitness effects of mutations fixed during virus adaptation to different ecotypes of a new host. *Philos Trans R Soc B Biol Sci* **370**, 20140292.
- Lapidot, M., Friedmann, M., Pilowsky, M., Ben-Joseph, R. & Cohen, S. (2001). Effect of host plant resistance to Tomato yellow leaf curl virus (TYLCV) on virus acquisition and transmission by its whitefly vector. *Phytopathology* **91**, 1209–1213.
- Larder, B. A. (1994). Interactions between drug resistance mutations in human immunodeficiency virus type 1 reverse transcriptase. *J Gen Virol* **75**, 951–957.
- Lefevre, P. & Moriones, E. (2015). Recombination as a motor of host switches and virus emergence: geminiviruses as case studies. *Curr Opin Virol* **10**, 14–19.
- Michelson, I., Zamir, D. & Czosnek, H. (1994). Accumulation and translocation of Tomato yellow Leaf curl virus (TYLCV) in a *Lycopersicon esculentum* breeding line containing the *L. chilense* TYLCV tolerance gene *Ty-1*. *Phytopathology* **84**, 928–933.
- Monci, F., Sánchez-Campos, S., Navas-Castillo, J. & Moriones, E. (2002). A natural recombinant between the geminiviruses Tomato yellow leaf curl Sardinia virus and Tomato yellow leaf curl virus exhibits a novel pathogenic phenotype and is becoming prevalent in Spanish populations. *Virology* **303**, 317–326.
- Moudy, R. M., Meola, M. A., Morin, L. L., Ebel, G. D. & Kramer, L. D. (2007). A newly emergent genotype of West Nile virus is transmitted earlier and more efficiently by *Culex* mosquitoes. *Am J Trop Med Hyg* **77**, 365–370.
- Pérez de Castro, A., Díez, M. J. & Nuez, F. (2005). Evaluation of breeding tomato lines partially resistant to *Tomato yellow leaf curl Sardinia virus* and *Tomato yellow leaf curl virus* derived from *Lycopersicon chilense*. *Can J Plant Pathol* **27**, 268–275.
- Pérfarres, F., Thébaud, G., Lefevre, P., Chiroleu, F., Rimbaud, L., Hoareau, M., Reynaud, B. & Lett, J.-M. (2014). Frequency-dependent assistance as a way out of competitive exclusion between two strains of an emerging virus. *Proc Biol Sci* **281**, 20133374.
- R_Development_Core_Team. (2010). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Richman, D. D., Havlir, D., Corbeil, J., Looney, D., Ignacio, C., Spector, S. A., Sullivan, J., Cheeseman, S., Barringer, K. & other authors (1994). Nevirapine resistance mutations of human immunodeficiency virus type 1 selected during therapy. *J Virol* **68**, 1660–1666.
- Rodríguez-Negrete, E. A., Carrillo-Tripp, J. & Rivera-Bustamante, R. F. (2009). RNA silencing against geminivirus: complementary action of post-transcriptional gene silencing and transcriptional gene silencing in host recovery. *J Virol* **83**, 1332–1340.
- Sánchez-Campos, S., Navas-Castillo, J., Camero, R., Soria, C., Díaz, J. A. & Moriones, E. (1999). Displacement of tomato yellow leaf curl virus (TYLCV)-Sr by TYLCV-Is in tomato epidemics in Spain. *Phytopathology* **89**, 1038–1043.
- Sahu, P. P., Sharma, N., Puranik, S. & Prasad, M. (2014). Post-transcriptional and epigenetic arms of RNA silencing: a defense machinery of naturally tolerant tomato plant against Tomato leaf curl New Delhi virus. *Plant Mol Biol Report* **32**, 1015–1029.
- Shi, M., Holmes, E. C., Brar, M. S. & Leung, F. C.-C. (2013). Recombination is associated with an outbreak of novel highly pathogenic porcine reproductive and respiratory syndrome viruses in China. *J Virol* **87**, 10904–10907.
- Tahiri, A., Sekkat, A., Bennani, A., Granier, M., Delvare, G. & Peterschmitt, M. (2006). Distribution of tomato-infecting begomoviruses and *Bemisia tabaci* biotypes in Morocco. *Ann Appl Biol* **149**, 175–186.

- Tahiri, A., Halkett, F., Granier, M., Gueguen, G. & Peterschmitt, M. (2013).** Evidence of gene flow between sympatric populations of the Middle East-Asia Minor 1 and Mediterranean putative species of *Bemisia tabaci*. *Ecol Evol* **3**, 2619–2633.
- Tomas, N., Zwart, M. P., Poulain, M. & Elena, S. F. (2014).** Estimation of the *in vivo* recombination rate for a plant RNA virus. *J Gen Virol* **95**, 724–732.
- Urbino, C., Gutiérrez, S., Antolik, A., Bouazza, N., Doumayrou, J., Granier, M., Martin, D. P. & Peterschmitt, M. (2013).** Within-host dynamics of the emergence of Tomato yellow leaf curl virus recombinants. *PLoS One* **8**, e58375.
- Vu, T. T., Holmes, E. C., Duong, V., Nguyen, T. Q., Tran, T. H., Quail, M., Churcher, C., Parkhill, J., Cardoso, J. & other authors (2010).** Emergence of the Asian 1 genotype of dengue virus serotype 2 in Viet Nam: in vivo fitness advantage and lineage replacement in South-East Asia. *PLoS Negl Trop Dis* **4**, e757.
- Vuillaume, F., Thébaud, G., Urbino, C., Forfert, N., Granier, M., Froissart, R., Blanc, S. & Peterschmitt, M. (2011).** Distribution of the phenotypic effects of random homologous recombination between two virus species. *PLoS Pathog* **7**, e1002028.
- Wain-Hobson, S., Renoux-Elbé, C., Vartanian, J. P. & Meyerhans, A. (2003).** Network analysis of human and simian immunodeficiency virus sequence sets reveals massive recombination resulting in shorter pathways. *J Gen Virol* **84**, 885–895.
- Worobey, M. & Holmes, E. C. (1999).** Evolutionary aspects of recombination in RNA viruses. *J Gen Virol* **80**, 2535–2543.
- Yadav, R. K. & Chattopadhyay, D. (2011).** Enhanced viral intergenic region-specific short interfering RNA accumulation and DNA methylation correlates with resistance against a geminivirus. *Mol Plant Microbe Interact* **24**, 1189–1197.
- Yang, X., Wang, Y., Guo, W., Xie, Y., Xie, Q., Fan, L. & Zhou, X. (2011).** Characterization of small interfering RNAs derived from the geminivirus/betasatellite complex using deep sequencing. *PLoS One* **6**, e16928.
- Zhou, X., Liu, Y., Calvert, L., Munoz, C., Otim-Nape, G. W., Robinson, D. J. & Harrison, B. D. (1997).** Evidence that DNA-A of a geminivirus associated with severe cassava mosaic disease in Uganda has arisen by interspecific recombination. *J Gen Virol* **78**, 2101–2111.