Review

Correspondence

Stephanie M. Rainey stephanie.graham@glasgow. ac.uk Isabelle Dietrich isabelle.dietrich@glasgow.ac.uk

Understanding the *Wolbachia*-mediated inhibition of arboviruses in mosquitoes: progress and challenges

Stephanie M. Rainey, Pranav Shah, Alain Kohl and Isabelle Dietrich

MRC - University of Glasgow Centre for Virus Research, 8 Church Street, Glasgow G11 5JR, UK

Arthropod-borne viruses (arboviruses) pose a considerable threat to human and animal health, yet effective control measures have proven difficult to implement, and novel means of controlling their replication in arthropod vectors, such as mosquitoes, are urgently required. One of the most exciting approaches to emerge from research on arthropods is the use of the endosymbiotic intracellular bacterium *Wolbachia* to control arbovirus transmission from mosquito to vertebrate. These α -proteobacteria propagate through insects, in part through modulation of host reproduction, thus ensuring spread through species and maintenance in nature. Since it was discovered that *Wolbachia* endosymbiosis inhibits insect virus replication in *Drosophila* species, these bacteria have also been shown to inhibit arbovirus replication and spread in mosquitoes. Importantly, it is not clear how these antiviral effects are mediated. This review will summarize recent work and discuss determinants of antiviral effectiveness that may differ between individual *Wolbachia*/vector/arbovirus interactions. We will also discuss the application of this approach to field settings and the associated risks.

Introduction

Arthropod-borne viruses (arboviruses) are transmitted in nature between susceptible vertebrate hosts and bloodfeeding arthropod vectors, such as mosquitoes, sandflies, midges and ticks. Upon infection with arboviruses, vertebrate hosts can develop a range of symptoms from mild to severe, including encephalitis and haemorrhagic fever. Arthropod vectors acquire a lifelong, generally asymptomatic infection through feeding on viraemic vertebrates. Following the initial replication in midgut or hindgut epithelial cells, virus is released into the haemocoel from where it disseminates to other tissues, most importantly the salivary glands and ducts. The transmission cycle is completed when arboviruses present in the arthropod's saliva are passed to a vertebrate host during a subsequent bloodmeal. Arbovirus distribution is dependent on the presence of vector species in a given area. Changes in climate and vector distribution, and an increase in urbanization, human travel and livestock movements, can all potentially impact on arbovirus transmission (Weaver & Reisen, 2010). Most arboviruses are grouped into four RNA virus families, and include the positive strand Togaviridae and Flaviviridae, the negative strand segmented Bunyaviridae and the dsRNA segmented Reoviridae (Weaver & Reisen, 2010).

Control of arboviral disease may be achieved by a number of measures, such as vaccination or the development of antiviral drugs; however, these are often not available and/or prove difficult to develop. For example, there is no effective vaccine for dengue virus (DENV; *Flaviviridae*), which impacts significantly on human health worldwide (Halstead, 2012) or emerging arboviruses, such as the alphavirus chikungunya (CHIKV; *Togaviridae*).

Indeed, preventing transmission from arthropod vector to vertebrate, and thus preventing infection altogether, may present a valuable alternative in combating arboviral disease. Vector eradication programmes form part of such strategies. In the past, sterile mosquitoes have been released, and more recently, genetic modification techniques have been used successfully to control mosquito populations (Alphey *et al.*, 2013; Black *et al.*, 2011; Lacroix *et al.*, 2012; Wilke & Marrelli, 2012). However, because one approach alone is unlikely to be sufficient and/or always applicable, novel strategies that work at the level of vector control are required.

In this context, current efforts to manipulate transmission of arboviruses by mosquitoes through the use of the bacterium *Wolbachia* merit greater attention. As RNA virus replication is inhibited by these endosymbionts in the model arthropod *Drosophila melanogaster* (Hedges *et al.*, 2008; Teixeira *et al.*, 2008), *Wolbachia* has also been used to control arbovirus replication in mosquitoes and their transmission to vertebrates.

Here we summarize recent research focusing on *Wolbachia*/ mosquito/arbovirus interactions and discuss progress into the mechanisms by which these endosymbionts confer antiviral resistance to their hosts, as well as perspectives for further research.

Wolbachia and reproductive phenotypes

Wolbachia is a genus of intracellular α -proteobacteria (family *Rickettsiaceae*), which is thought to infect 40–75% of all arthropod species and several nematodes (Hilgenboecker *et al.*, 2008; Zug & Hammerstein, 2012). First discovered in the mosquito *Culex pipiens* (Hertig & Wolbach, 1924), these bacteria have now been found in a number of vector mosquito species, although not the major arbovirus vector *Aedes aegypti* (Kittayapong *et al.*, 2000; Sinkins, 2004).

Wolbachia has been extensively studied because of its ability to significantly alter the biology of the host in the symbiotic relationship. These alterations are thought to dramatically increase the spread of the bacteria through insect populations and include male killing, parthenogenesis, feminization and cytoplasmic incompatibility (CI) (Werren *et al.*, 2008). In mosquito populations, *Wolbachia* is spread and maintained through unidirectional and bidirectional CI (Kambhampati *et al.*, 1993; Sinkins, 2004; Yen & Barr, 1973), and our understanding of this process is vital to successful introduction of *Wolbachia*-infected mosquitoes into wild populations (as discussed in the final section of this review). The process of CI is explained below and summarized in Fig. 1.

Alterations in spermatogenesis during male development result in the failure of *Wolbachia*-infected males to produce offspring when mating with either uninfected females (unidirectional CI) or females infected with a different strain of *Wolbachia* (bidirectional CI) (Fig. 1). The altered sperm results in late entry of the male pronucleus into mitosis. Thus, the parental chromosomes fail to align properly and haploid embryos are produced. The presence of compatible *Wolbachia* in fertilized eggs, through vertical transmission, rescues CI (Landmann *et al.*, 2009; Stouthamer *et al.*, 1999; Tram *et al.*, 2006; Tram & Sullivan, 2002). Importantly, females infected with *Wolbachia* can mate successfully with uninfected males, giving them an evolutionary advantage over uninfected females, as they produce viable offspring with both infected and uninfected males (Turelli & Hoffmann, 1991, 1995; Yen & Barr, 1971).

As early as the 1960s, mosquitoes that exhibited CI were considered a viable option for the control of disease vectors. One study reported the large scale release of Wolbachiainfected C. pipiens into wild populations in order to eliminate this population from a large urban area (Laven, 1967). More recently, a virulent strain of Wolbachia found in laboratory strains of D. melanogaster, wMelPop, was cited as a possible mechanism for the control of mosquito populations and consequently the spread of arboviruses through transinfection (see below) (Brownstein et al., 2003; Yeap et al., 2011). This particular strain of Wolbachia is known to infect Drosophila at a high density and reduce lifespan. It has since been shown that it is possible to transinfect Aedes albopictus and Ae. aegypti populations with wMelPop and other Wolbachia strains, and for them to be stably maintained in populations (Blagrove et al., 2013; McMeniman et al., 2009; Xi et al., 2005). The potential of Wolbachia to control arbovirus spread, however, was not fully realized until 2008 when two groups reported that Wolbachia infection in D. melanogaster led to increased protection against insect RNA viruses, suggesting that Wolbachia was able to control viral replication in infected flies (Hedges



Fig. 1. Cl induced by *Wolbachia* infection. Cl is presented in two forms in insects. (a) The process by which bidirectional Cl takes place. Here, males and females infected with two different strains of *Wolbachia* are unable to produce viable offspring. (b) The process of unidirectional Cl, where infected females are able to successfully mate with both uninfected males and males infected with the same strain of *Wolbachia*.

et al., 2008; Teixeira et al., 2008). The remainder of this review will discuss this in more detail.

Wolbachia and antiviral activity in insects

Studies in the model Drosophila

Initial studies carried out in *D. melanogaster* concluded that the *Wolbachia* strains *w*MelCS and *w*MelPop increased fly longevity upon infection with the RNA viruses *Drosophila* C virus (DCV; *Dicistroviridae*), Flock House virus (FHV; *Nodaviridae*) and cricket paralysis virus (CrPV; *Dicistroviridae*), but not the DNA virus insect iridescent virus 6 (IIV-6; *Iridoviridae*) (Hedges *et al.*, 2008; Teixeira *et al.*, 2008). Moreover, viral titres were significantly reduced in the presence of *Wolbachia* in the case of DCV and nora virus (NoraV; a potentially novel family within the *Picornavirales*).

Further studies in *Drosophila simulans*, which are naturally infected with multiple *Wolbachia* strains, showed that only those *Wolbachia* strains that are present in high densities (*w*Mel, *w*Ri and *w*Au), conferred antiviral resistance to DCV and FHV. In DCV-infected *D. simulans*, *Wolbachia*-mediated protection did not directly correlate with reduced viral copy numbers (Osborne *et al.*, 2009). For example,

wRi significantly increased lifespan upon infection with DCV compared to controls; however, there was no significant decrease in virus accumulation. Conversely, Teixeira et al. (2008) and Hedges et al. (2008) showed that DCV titres in D. melanogaster were strongly reduced by Wolbachia infection (Hedges et al., 2008; Teixeira et al., 2008). During FHV infection, virus titre was unaffected but virusinduced pathology was decreased, resulting in increased life spans of infected flies (Teixeira et al., 2008). A recent study also demonstrated that replication of the dsRNA virus bluetongue virus (BTV; Reoviridae) in D. melanogaster is inhibited in the presence of Wolbachia (Shaw et al., 2012). Taken together these data suggest that reduction of viral titres is not the only mechanism responsible for Wolbachiamediated protection. These findings in the model organism Drosophila have led to considerable interest in the field of human-pathogenic mosquito-borne viruses. The key findings are summarized in Table 1.

Mosquitoes and Wolbachia-mediated antiviral activity

Aedes and Culex species are the major arthropod vectors for DENV, West Nile virus (WNV) and yellow fever virus (YFV) (all *Flaviviridae*), as well as CHIKV (*Togaviridae*) (Weaver & Reisen, 2010). Among the major arbovirus

Species	Wolbachia strain	Nature of host- <i>Wolbachia</i> association	Virus	Antiviral effect exerted by <i>Wolbachia</i>	Reference
Drosophila melanogaster	wMelCS	Natural	DCV	Increased lifespan of infected flies; reduced virus proliferation	(Teixeira et al., 2008)
			FHV*	Increased lifespan of infected flies; no effect of virus replication	
			NoraV	Reduced virus proliferation	
			IIV-6	Decreased lifespan of infected flies; increased virus proliferation	
			DCV, FHV*, CrPV	Increased lifespan of infected flies; reduced virus proliferation	(Hedges <i>et al.</i> , 2008)
	wMelPop	Laboratory strain	DCV, FHV*, CrPV	Increased lifespan of infected flies; reduced virus proliferation	(Hedges <i>et al.</i> , 2008)
	wMel	Natural	CHIKV*, LACV*	No effect	(Glaser & Meola, 2010)
			WNV*	Reduced virus proliferation	(Glaser & Meola, 2010)
			BTV*	Reduced virus proliferation	(Shaw et al., 2012)
Drosophila simulans	wMel	Transinfection	DCV	Increased lifespan of infected flies; reduced virus proliferation	(Osborne <i>et al.</i> , 2009)
	wAu	Natural	DCV	Increased lifespan of infected flies; reduced virus proliferation	(Osborne <i>et al.</i> , 2009)
			FHV*	Increased lifespan of infected flies;	(Osborne et al., 2009)
	wRi	Natural	DCV	Increased lifespan of infected flies	(Osborne et al., 2009)
			FHV*	Increased lifespan of infected flies	(Osborne et al., 2009)
	wHa	Natural	DCV, FHV*	No effect	(Osborne et al., 2009)
	wNo	Natural	DCV, FHV*	No effect	(Osborne et al., 2009)

BTV, Bluetongue virus; CHIKV, chikungunya virus; CrPV, cricket paralysis virus; DCV, *Drosophila* C virus; FHV, Flock House virus; IIV-6, insect iridescent virus 6; LACV, La Crosse virus; NoraV, nora virus; WNV, West Nile virus.

vectors, *Ae. albopictus*, *Aedes bromeliae* and members of the *C. pipiens* complex are naturally infected with *Wolbachia* endosymbionts (*Ae. albopictus* harbours wAlbA and wAlbB, and *C. pipiens wPip*) (Armbruster *et al.*, 2003; Hertig & Wolbach, 1924; Kittayapong *et al.*, 2000; Osei-Poku *et al.*, 2012; Sinkins *et al.*, 1995), whereas *Ae. aegypti* mosquitoes lack this association (Table 2).

Unlike the above-mentioned native Drosophila/Wolbachia interactions, the ability of homologous Wolbachia strains to confer resistance in mosquitoes is limited by tissue density and distribution. For example, wAlbA and wAlbB did not inhibit DENV replication to significant levels in Ae. albopictus mosquitoes (Bian et al., 2010), but reduced viral infection of the salivary glands and thus may limit transmission (Mousson et al., 2012). Furthermore, the wAlbA and wAlbB infected Ae. albopictus cell line Aa23 showed decreased DENV titres compared to Wolbachiacured controls. Notably this cell line has a significantly higher Wolbachia density than somatic tissues of Ae. albopictus, which may explain its restrictive phenotype (Lu et al., 2012). A study of the interaction between CHIKV and wAlbA and wAlbB in Ae. albopictus mosquitoes showed no decrease in viral titres in the presence of Wolbachia compared to controls. Viral titres in Wolbachia-free mosquitoes were extremely variable compared to those harbouring Wolbachia, suggesting some level of interaction (Mousson et al., 2010). Lastly, homologous wPip infections in Culex quinquefasciatus resulted in reduced WNV titres and transmission rates (Glaser & Meola, 2010). Given that *Wolbachia* confers resistance to RNA viruses in *Drosophila* (Hedges *et al.*, 2008; Teixeira *et al.*, 2008), and that *Wolbachia* strains can be stably introduced and maintained in heterologous host mosquitoes (Blagrove *et al.*, 2013; McMeniman *et al.*, 2009; Xi *et al.*, 2005), multiple studies have investigated whether *Wolbachia* transinfection into different mosquito species would result in vectors refractory to infection with important viral pathogens.

Transinfection and resulting phenotypes

Transinfection of mosquitoes with heterologous Wolbachia strains is by no means trivial, as not all hosts are equally permissive and Wolbachia strains differ in their ability to stably transfect certain new host species (Werren et al., 2008). Most likely as a consequence of evolutionary adaptation to their hosts, naturally occurring Wolbachia strains differ in their genome organization and in the reproductive phenotypes they cause (reviewed by Serbus et al., 2008; Werren et al., 2008). In nature, Wolbachia cross-species transmission events occur frequently (Haine et al., 2005; Heath et al., 1999; Huigens et al., 2000, 2004; Kraaijeveld et al., 2011; Vavre et al., 1999; Werren et al., 1995) and are thought to result in a loss of infection or, if infection is established, in a temporary increase in virulence, caused either by competition for resources or by disproportional host immune activation (Le Clec'h et al., 2012; Lipsitch

Mosquito host	<i>Wolbachia</i> strain	Nature of host- <i>Wolbachia</i> association	Virus	Antiviral effect exerted by <i>Wolbachia</i>	Reference
Aedes aegypti	wAlbB	Transinfection	DENV	Reduced virus proliferation	Bian et al. (2010)
	wMel	Transinfection	DENV	Blockage of viral proliferation	Walker et al. (2011)
			WNV	No effect	Hussain et al. (2013)
			CHIKV, YFV	Reduced virus proliferation	van den Hurk et al. (2012)
	wMelPop	Transinfection	DENV, CHIKV	Reduced virus proliferation	Moreira et al. (2009)
			YFV	Reduced virus proliferation	van den Hurk et al. (2012)
			WNV	Reduced virus proliferation	Hussain et al. (2013)
Aedes albopictus	wAlbA and wAlbB	Natural	CHIKV	No effect	Mousson et al. (2010)
			DENV	No effect on virus replication in midgut; reduced virus dissemination and transmission	
			DENV	No effect	Lu et al. (2012)
	<i>w</i> Mel (<i>w</i> AlbA and <i>w</i> AlbB present)	Transinfection	CHIKV	Reduced virus transmission	Blagrove et al. (2013)
Aedes	wPolA	Natural	DENV	No effect	Bian et al. (2013b)
polynesiensis	wAlbB	Transinfection	DENV	Reduced virus proliferation	Bian et al. (2013b)
Culex quinquefasciatus	wPip	Natural	WNV	Reduced virus proliferation	Glaser & Meola (2010)
Armigeres subalbatus	Not subgrouped	Natural	JEV	No effect in salivary gland cells	Tsai et al. (2006)

Table 2. Antiviral effect of Wolbachia in different mosquito/arbovirus associations

CHIKV, Chikungunya virus; DENV, dengue virus; JEV, Japanese encephalitis virus; WNV, West Nile virus; YFV, yellow fever virus.

& Moxon, 1997). In order to stably introduce the D. melanogaster Wolbachia strain wMelPop into Ae. aegypti, McMeniman et al. (2008) first had to pre-adapt wMelPop to an Ae. albopictus-derived cell line by long-term serial passage (McMeniman et al., 2008). After about 2.5 years in culture the pre-adapted strain was transferred into Ae. aegypti and Anopheles gambiae cell lines and passaged further. The resulting Wolbachia strain (named wMelPop-CLA) was then reintroduced into D. melanogaster cured of wMelPop, where it showed reduced virulence and ability to induce CI compared to wMelPop (McMeniman et al., 2008). In Ae. aegypti, wMelPop-CLA induced life-shortening and CI (McMeniman et al., 2009). Further to this, several species of Wolbachia from both Drosophila and mosquito species have been transinfected into different hosts, leading to different levels of antiviral activity (summarized in Table 2). For example, in wMel-transinfected Ae. aegypti, WNV was not inhibited, whereas the presence of the *w*MelPop strain of Wolbachia significantly reduced WNV replication (Hussain et al., 2013). Similarly, wMelPop-CLA, and to a lesser extent wMel, was able to diminish YFV replication and dissemination in transinfected Ae. aegypti (van den Hurk et al., 2012). In the same study it has been shown that wMel can only inhibit infection and dissemination of CHIKV in transinfected Ae. aegypti upon viraemic bloodmeal, but not upon intrathoracic inoculation (van den Hurk et al., 2012). More recently, wMel transinfection in Ae. albopictus was found to prevent CHIKV dissemination to the salivary glands and thus possibly viral transmission (Blagrove et al., 2013).

Further studies have shown that transinfection of *Ae. aegypti* with the *D. melanogaster w*MelPop-CLA strain of *Wolbachia* negatively affects mosquito survival and strongly inhibits the replication of DENV when compared to tetracycline-treated (cured of *Wolbachia* infection) control or wild-type mosquitoes, irrespective of whether mosquitoes were blood fed or whether virus was injected intrathoracically (Moreira *et al.*, 2009). *Wolbachia* also reduced DENV dissemination to the thorax and head of mosquitoes, and hence possibly transmission (Bian *et al.*, 2010; Moreira *et al.*, 2009). *Ae. aegypti* transinfected with the *Ae. albopictus Wolbachia* strain *w*AlbB also showed increased mosquito longevity upon infection with DENV and suppressed viral replication (Bian *et al.*, 2010).

It would appear that the *Wolbachia*-induced virus refractory phenotype is dependent on the combination of *Wolbachia* strain, virus and host (genetic or other) factors. Understanding these differences is key to a successful vector control protocol. Several mechanisms have been proposed to explain why *Wolbachia* inhibits arbovirus transmission effectively in some scenarios but not in others. These will be discussed below.

Determinants of *Wolbachia* viral interference in arthropod hosts

It is not known how *Wolbachia* endosymbionts confer antiviral resistance to their hosts, and in order to develop intervention strategies based on *Wolbachia*-mediated viral interference it is crucial to elucidate the underlying mechanisms. An important question to address is whether *Wolbachia* interferes with viral replication directly or whether it increases vector resistance to viral infection by other mechanisms. Furthermore, it is important to elucidate whether the *Wolbachia*-mediated effect is cell autonomous or systemic.

Wolbachia density and competition for host cell resources

Density. Early studies found a positive correlation between intracellular endosymbiont density and antiviral effect conferred (Lu et al., 2012; Osborne et al., 2009, 2012). Different Wolbachia strains are known to infect vector species at variable densities and to display different tissue distributions (Dobson et al., 1999; Dutton & Sinkins, 2004; Miller & Riegler, 2006). In its native host Ae. albopictus, wAlbB is found both in reproductive tissues (ovaries, testes) and somatic tissues (haemolymph, midgut, muscle, head) (Dobson et al., 1999). In somatic tissues, wAlbB levels can vary significantly between Ae. albopictus strains (Dobson et al., 1999). In C. pipiens, Wolbachia densities can vary up to 100-fold between individual mosquitoes and also between populations, which could lead to differences in vector competence (Berticat et al., 2002; Echaubard et al., 2010; Glaser & Meola, 2010). In heterologous hosts, however, Wolbachia usually grows to higher densities than in native hosts, which may explain why antiviral effects are more often observed in heterologous hosts compared to native host species (Glaser & Meola, 2010). Examples of Wolbachia transinfections that resulted in increased bacterial densities and antiviral resistance compared to original vector species include infection of Ae. aegypti (Bian et al., 2010) or Aedes polynesiensis (Bian et al., 2013b) with wAlbB. An exception is wMelPop, which grows to high titres in D. melanogaster and leads to a reduced life span of the host and a reduction in viral replication (Hedges et al., 2008; Min & Benzer, 1997).

It has been hypothesized that protection is dependent on whether *Wolbachia*-harbouring tissues and viral target cells or tissues co-localize (Bian *et al.*, 2010; Moreira *et al.*, 2009). Interestingly, in *C. pipiens Wolbachia* is less abundant in the midgut than in other somatic tissues (Dobson *et al.*, 1999) and may not be present to limit DENV replication in midgut epithelial cells (Glaser & Meola, 2010). Density may also be important if it is simply a case of competition for space and/or cellular resources. Indeed there is little evidence of *Wolbachia* and virus being present together in the same cell/tissue when either is at a high density (Frentiu *et al.*, 2010; Moreira *et al.*, 2009).

Autophagy. One mechanism that is thought to control the density of *Wolbachia* is autophagy. Infection of C6/36 cells (an *Ae. albopictus*-derived cell line) with *w*AlbB showed that *Wolbachia* not only induced the autophagy pathway

but also manipulated it in order to ensure survival (Voronin *et al.*, 2012). This was also true for a naturally infected *D. melanogaster* line (*w*MelPop) (Voronin *et al.*, 2012). Since both alphaviruses and flaviviruses are thought to require the autophagy pathway in order to replicate (Krejbich-Trotot *et al.*, 2011; Lee *et al.*, 2008), this cellular function may be a site of *Wolbachia* interference with virus replication. Further work is clearly required to confirm this proposal.

Iron metabolism and cholesterol. If the *Wolbachia*mediated antiviral effect is cell autonomous, then one explanation for reduced virus replication may be competition for resources between *Wolbachia*, viruses and the host cell. For example, iron homeostasis has been shown to be regulated by *Wolbachia* (Kremer *et al.*, 2009), with transcriptomic data suggesting that iron-binding proteins, transferrin and ferritin, were upregulated in the presence of *w*Mel and *w*MelPop-CLA in *Ae. aegypti* (Rancès *et al.*, 2012). Data have further indicated that transferrin was downregulated upon infection with DENV and CHIKV (Tchankouo-Nguetcheu *et al.*, 2010).

In addition to iron homeostasis, the cholesterol metabolism needs to be tightly regulated to enable viral replication and bacterial growth. Insect cells are cholesterol auxotrophs (Clayton, 1964), and can be depleted of cholesterol and other lipids by growth in delipidated serum (Hafer *et al.*, 2009; Silberkang *et al.*, 1983); however, they will incorporate low levels of cholesterol from serum (Mitsuhashi *et al.*, 1983). *Wolbachia* is unable to synthesize cholesterol *de novo* and thus its replication is cholesteroldependent (Lin & Rikihisa, 2003). Further, inside host cells individual bacteria are contained within cholesterol-containing Golgi-related vesicles (see also Fig. 2) (Cho *et al.*, 2011). Cholesterol depletion of host cells by *Wolbachia* could also interfere with virus replication. A recent study has examined the involvement of cholesterol in *Wolbachia*-mediated DCV interference in *D. melanogaster* (Caragata *et al.*, 2013). It was shown that cholesterol enrichment of fly diets reversed the *Wolbachia*-mediated DCV interference effect and the increase in *D. melanogaster* life span in a dose-dependent manner. Importantly, dietary cholesterol levels did not affect *Wolbachia* densities in *w*MelCS-infected flies and did not affect DCV infection levels in *Wolbachia*-free flies (Caragata *et al.*, 2013). For *w*MelPop-infected flies there was a significant effect of diet on *Wolbachia* density; however, this did not affect pathogen blocking (Caragata *et al.*, 2013).

Both mosquito-borne alphaviruses and flaviviruses have been shown to rely on host cell lipids for their uptake, replication, virion assembly and infectivity. Sindbis virus (SINV; Togaviridae) and Semliki Forest virus (SFV; Togaviridae) are endocytically taken up into host cells, followed by cholesterol and sphingolipid-dependent fusion of the viral and host membranes at low pH (Kielian, 1995; Lu et al., 1999; Smit et al., 1999). Their replication takes place in replication complexes on intracellular endosomederived membranes. In insect cells, virion assembly takes place in cytoplasmic vesicles that fuse with the plasma membrane in a cholesterol-dependent manner to release virus into the medium (Brown & Condreay, 1986). Therefore, the lipid content of viral particles resembles that of the producer cells (Hafer et al., 2009). Several studies have shown that alphaviruses can grow to high titres in insect cells although their lipid content is generally low (Hafer et al., 2009; Mitsuhashi et al., 1983; Tsetsarkin et al., 2011). Cholesterol depletion of C6/36 cells blocked SFV endosome membrane fusion, replication and exit (Chatterjee et al., 2000; Marquardt et al., 1993; Phalen & Kielian, 1991). A significant reduction in the production of infectious SINV and SFV particles upon delipidation of C6/36 cells has been reported (Hafer et al., 2009; Lu et al., 1999; Marquardt et al., 1993; Phalen & Kielian, 1991). It can be assumed that the lower infectivity of these particles was caused by an



Fig. 2. Transmission electron micrographs of *Drosophila* JW18 cell line infected with *Wolbachia*. (a) Low magnification image of a whole cell depicting the distribution of *Wolbachia* (W) in *Drosophila* JW18 cells (Serbus *et al.*, 2012), with respect to the nucleus (N) and vacuoles (V). (b) Higher magnification of the boxed area of the cell in (a) showing a pair of *Wolbachia* cells in the same vacuole, presumably at the end of the process of division. Bars, 1 μm (a) and 0.5 μm (b).

altered lipid composition, which resulted in a loss of stability and structural fragility of the produced virions. Interestingly, in addition to cholesterol other lipids seem to be essential for virion infectivity (Hafer *et al.*, 2009).

Similar to alphaviruses, flavivirus replication relies on a tight regulation of host cell cholesterol biosynthesis (Carro & Damonte, 2013; Rothwell et al., 2009). Although DENV was found to be insensitive to plasma membrane cholesterol depletion in C6/36 cells (Acosta et al., 2008; Mosso et al., 2008; Umashankar et al., 2008), extraction of cholesterol from viral envelopes reduced DENV infectivity for C6/36 cells (Acosta et al., 2009). The differential dependence on host cell membrane cholesterol content seen between members of the families Togaviridae and Flaviviridae may be explained by the fact that the former viruses bud through the host cell plasma membrane so that the lipid composition of their envelopes resembles that of the cell membrane (Hafer et al., 2009; Sousa et al., 2011). In contrast, the origin of the flavivirus envelope is the endoplasmic reticulum (Welsch et al., 2009), which contains only a small proportion of the total cell cholesterol (Lange et al., 1999). Given the crucial role for cholesterol and other lipids in viral life cycles, it is not surprising that another flavivirus, WNV, has been shown to modulate host cell cholesterol levels by upregulating cholesterol biosynthesis and redistributing it to sites of virus replication (Mackenzie et al., 2007). These cellular modifications are necessary for WNV replication to take place. In summary, these early experiments relating to Wolbachia, cholesterol and antiviral activity are intriguing and warrant further investigation as it is clear that cholesterol is important both for virus replication and Wolbachia growth.

Immune signalling in arthropod/Wolbachia interactions

As Wolbachia is now known to confer resistance to a broad range of RNA viruses, it could be assumed that the mechanism(s) behind antiviral activity would also be broad ranging. Innate immunity consists of a set of anti-microbial mechanisms largely conserved between insect species and is the first line of defence against invading microbes, including viruses (Bartholomay *et al.*, 2010; Christophides *et al.*, 2002; Lemaitre & Hoffmann, 2007; Waterhouse *et al.*, 2007). It is therefore plausible to suggest that Wolbachia may control viral infection by pre-activating the immune system, also known as immune priming. Much of our understanding of innate immunity in mosquitoes is influenced by key findings made in *Drosophila* studies (Fragkoudis *et al.*, 2003; Wong *et al.*, 2011).

Toll, Imd and JAK-STAT

Among a number of mechanisms that limit bacterial infections, three major signalling pathways – Toll, Imd and JAK-STAT – control innate immunity. The Toll pathway

is initiated upon infection with fungi or Gram-positive bacteria, leading to transcriptional upregulation of a specific subset of antimicrobial peptides (AMPs) and several other immune genes (Lemaitre & Hoffmann, 2007). The Toll pathway has also been implicated in antiviral immunity. For example, the pathway is activated by the presence of the dsRNA virus Drosophila X virus (DXV; Birnaviridae) in Drosophila (Zambon et al., 2005), and has been implicated in the control of DENV and WNV in mosquito studies (Smartt et al., 2009; Xi et al., 2008). Stimulation with Gramnegative bacteria activates the Imd pathway, and induces immune genes and Imd-specific AMPs (Lemaitre & Hoffmann, 2007). Several mutants in the Drosophila Imd pathway also exhibit higher CrPV titres compared to wildtype flies (Costa et al., 2009). Studies in An. gambiae suggest the differential expression of Imd pathway genes in response to o'nyong'nyong virus (ONNV; Togaviridae); however, it would only appear to have a minor role in that particular host-virus interaction (Waldock et al., 2012). A third pathway known to be involved in both bacterial and viral infection is the JAK-STAT pathway. DCV, for example, activates the JAK-STAT pathway in Drosophila resulting in the upregulation of genes such as vir1 (Dostert et al., 2005). Studies in Ae. aegypti have demonstrated that DENV infection is controlled by JAK-STAT signalling (Souza-Neto et al., 2009; Xi et al., 2008). Indeed, knockdown of key components of the JAK-STAT pathway resulted in significantly higher DENV viral titres. In the case of alphaviruses, a number of studies in Drosophila and mosquito cells suggest that the combination of Imd and/or JAK-STAT signalling, but not the Toll pathway, has antiviral activities (Avadhanula et al., 2009; Fragkoudis et al., 2008; Huang et al., 2013). Importantly, this shows that at least when it comes to antiviral immune signalling, differences between virus families, but also within virus families and vectors (SINV/SFV versus ONNV), exist.

Early studies in Ae. aegypti mosquitoes transinfected with several different strains of Wolbachia indicated that immune priming may be involved in antiviral activity. For example, the transinfection of wMelPop-CLA into Ae. aegypti resulted in the upregulation of immune genes, including several involved in the Toll and Imd pathways (Kambris et al., 2010; Rancès et al., 2012). Further transcriptional analysis of transinfected Ae. aegypti (wAlbB) showed significant upregulation of several Toll-dependent genes, in particular the AMP-encoding gene defensin (Bian et al., 2010). These studies are, however, hampered by the initial transinfection of heterologous Wolbachia strains. Naturally infected D. melanogaster, D. simulans, and indeed, Ae. albopictus do not show this immune priming phenotype, and in the case of D. simulans and D. melanogaster (and to some extent Ae. albopictus) Wolbachia still confers resistance to several RNA viruses (Lu et al., 2012; Osborne et al., 2009; Rancès et al., 2012; Teixeira et al., 2008). Significantly, DENV infection is limited in both D. melanogaster and Ae. *aegypti*, which are infected with the strain *w*Mel, to similar degrees (Rancès et al., 2012). Therefore, it is unlikely that the immune priming seen in Ae. aegypti is key to Wolbachiamediated protection, as there is no such priming in D. melanogaster. It has been shown that there was neither an induction nor suppression of AMPs in either D. simulans or Ae. albopictus naturally infected with different Wolbachia strains (see Tables 1 and 2, respectively) and these insects are furthermore able to mount a normal response to bacterial infection (Bourtzis et al., 2000). Immune priming of the Imd or Toll pathways would also suggest that Wolbachia had the ability to protect against further bacterial infection; however, again this is not the case (Rottschaefer & Lazzaro, 2012). A more recent study has utilized the availability of Drosophila strains that are deficient in key components of the Imd and Toll pathway to conclude that neither a functional Toll nor an Imd pathway is required for the ability of Wolbachia to protect against DENV (Rancès et al., 2013).

The Toll pathway has also been shown to be induced by the production of reactive oxygen species (ROS) in Ae. aegypti mosquitoes transinfected with the heterologous Wolbachia strain wAlbB (Pan et al., 2012). The activation of the pathway in these mosquitoes leads to the production of the AMPs cecropin and defensin. Targeted reduction of these two AMPs resulted in an increase of DENV viral titres in these mosquitoes, leading to speculation that production of ROS and subsequent Toll activation may lead to Wolbachia-mediated antiviral activity. However, the corresponding experiment was not carried out in Wolbachiafree lines; therefore, the extent by which this activation leads to antiviral activity cannot be fully assessed (Pan et al., 2012). The broad activity of Wolbachia-mediated protection would argue against immune priming as the main mediator of antiviral activity, given the different susceptibilities between viruses to Toll, Imd and JAK-STAT pathways.

A recent study identified the phenoloxidase (PO) cascade as an important mosquito innate immune response to SFV (Rodriguez-Andres *et al.*, 2012). Interestingly, *Wolbachia* has been shown to increase melanization through the PO cascade in both homologous and heterologous *Wolbachia*/ host settings (Thomas *et al.*, 2011). It is likely that the PO cascade acts to protect against other viruses, and therefore the interaction with *Wolbachia* is intriguing and would merit further investigation.

Taken together, the studies described above suggest that immune priming is not central to the protection that *Wolbachia* confers. However, as transinfection of *Wolbachia* strains to heterologous hosts often leads to some level of immune priming, the additive effect of this cannot be ignored. It may be that the amount of antiviral protection present in released mosquitoes is vital to the sustainability of vector/arbovirus control. Studies suggest that as there is little or no immune priming present in homologous *Wolbachia*/ host settings, over time co-evolution would lead to a reduction in the immune response. Therefore, it is crucial to understand to what extent immune priming plays a role in antiviral activity in transinfected hosts.

Small RNA pathways in arthropod/*Wolbachia* interactions

Further to the immune signalling pathways discussed above, mosquitoes respond to viral infection through the sequence-specific small RNA breakdown pathways, RNA interference (RNAi). RNAi is considered to be the most important antiviral response in insects and is key to the control of arboviruses in mosquitoes (Blair, 2011). The exact mechanisms behind antiviral RNAi are beyond the reach of this review (see Donald et al., 2012 for a recent review). Nevertheless, as a mechanism that is known to successfully control a wide variety of arboviruses, it is important to ascertain if the RNAi response is involved in mediating the antiviral activity of Wolbachia. Several lines of evidence, however, suggest that it is not essential. Firstly, the Ae. albopictus cell line C6/36 does not contain a functional Dicer 2 protein, a key protein in the main antiviral RNAi response (Brackney et al., 2010; Scott et al., 2010). Transinfection of these cells with wMel still results in the inhibition of DENV, suggesting that a functioning exogenous RNAi response is not necessary for inhibition to occur (Frentiu et al., 2010). Additionally, Drosophila mutants defective for several key components of the antiviral RNAi pathway infected with Wolbachia still show resistance to WNV, DCV and FHV when compared to control flies (Glaser & Meola, 2010; Hedges et al., 2012). Taken together these results indicate that the main antiviral RNAi pathway is not essential for Wolbachia-mediated viral protection.

In addition to the RNAi pathway, the microRNA (miRNA) pathway has been implicated in the pathogen-blocking effect of Wolbachia (Hussain et al., 2011; Osei-Amo et al., 2012; Zhang et al., 2013). miRNAs are nonprotein-coding small RNAs that are involved in the regulation of cellular development, differentiation, apoptosis and immunity (Asgari, 2013; Donald et al., 2012). Recently, the expression of miRNAs in Ae. aegypti mosquitoes transinfected with wMelPop-CLA was studied (Hussain et al., 2011). The presence of Wolbachia has been shown to alter the expression of several insect miRNAs. The induction of one miRNA, aae-miR-2940, by Wolbachia is of particular interest to this review. aae-miR-2940 has been shown to downregulate the expression of the metalloprotease m41 ftsh (Hussain et al., 2011). Inhibition of this miRNA or silencing of the metalloprotease gene in mosquitoes reduced Wolbachia density (Hussain et al., 2011). Importantly, upregulation of aae-miR-2940 by Wolbachia has been found to inhibit DENV replication. This effect on DENV is mediated by the downregulation of the Ae. aegypti DNA (cytosine-5) methyltransferase gene AaDnmt2, a second target of aae-miR-2940 (Zhang et al., 2013). DNA cytosine methylation is essential for host defence, genome stability, gene regulation, organ differentiation and ageing. Reversely, overexpression of AaDnmt2 leads to a decrease in Wolbachia density, but an increase in DENV replication. In DENV-infected, Wolbachia-free mosquitoes, expression of AaDnmt2 is upregulated. In Wolbachia-free Ae. aegypti, aae-miR-2940 is, although at low levels, expressed so that AaDnmt2 levels are negligible (Zhang *et al.*, 2013). These findings point towards an important role of aae-miR-2940 in the maintenance of *Wolbachia* infection in *Ae. aegypti* as well as the resistance to DENV in *Wolbachia*-transinfected mosquitoes. Reduction in *AaDnmt2* expression in *Wolbachia*-infected cells could lead to hypomethylation of the mosquito genome and thereby regulate expression of methylation-sensitive host genes (Zhang *et al.*, 2013). In fact, a recent study has confirmed this idea (Ye *et al.*, 2013). In particular, expression of *Ae. aegypti* genes, with membrane functions, appear to be differentially regulated upon *Wolbachia* infection (Cho *et al.*, 2011; Ye *et al.*, 2013).

In summary, the studies discussed above suggest that the RNAi pathway is not essential for *Wolbachia*-mediated antiviral activity, but that the miRNA pathway could affect viral replication either directly, through up or down-regulation of genes involved in viral replication or antiviral activity, or indirectly, by altering the density of *Wolbachia*.

Applications, risks and future directions

Based on its pathogen-blocking effect Wolbachia introduction into wild mosquito populations has been suggested as a potential control measure for arboviruses. Previous introductions of genetically modified (GM) mosquitoes into wild populations have been met with public concern. However, Wolbachia is a naturally occurring symbiont of mosquitoes, and therefore infected vectors are not considered to be GM. Recent studies have also demonstrated that there is no antibody production against Wolbachia in human volunteers bitten by Wolbachia-infected mosquitoes, nor is there a transfer to the environment (be it mosquito predators, or soil, leaves, etc.) (Popovici et al., 2010), suggesting that Wolbachia is a safe alternative to GM mosquitoes. Nevertheless, public engagement and detailed regulations, such as those set out by Hoffmann et al. (2011), are key to public and scientific acceptance of such release experiments.

Release of *Wolbachia*-infected mosquitoes into wild mosquito populations, with the aim of fixation of the infection frequency (i.e. stable introduction of *Wolbachia* in previously non-infected vector populations), requires extensive knowledge of local and regional mosquito population densities and dispersion dynamics (Engelstädter & Telschow, 2009; Schofield, 2002). Further parameters to consider include the strength of the *Wolbachia*-induced CI effect, maternal inheritance rates and fitness costs or advantages (for example refractoriness to viral infection) to the vector inferred by *Wolbachia* infection (Vavre & Charlat, 2012). Mathematical modelling approaches are then employed to calculate the minimum number of *Wolbachia*-infected mosquitoes to be released (Hancock & Godfray, 2012; Hoffmann *et al.*, 2011; Hughes & Britton, 2013; Turelli, 2010; Walker *et al.*, 2011).

A key question is how sustainable a *Wolbachia*-mediated approach would be. In order to answer this question, we firstly need to distinguish between two mechanisms by which stable introduction of *Wolbachia* into wild populations can limit vector competence for pathogen transmission: (i) Wolbachia-mediated life-shortening of mosquitoes, which would interfere with viral transmission, and (ii) Wolbachia-induced vector refractoriness to infection, which would limit viral replication and transmission (Vavre & Charlat, 2012). Modelling of the epidemiology of Wolbachia in host populations has suggested that vector/ Wolbachia associations would be more sustainable in the field using the former approach as host resistance to the bacterium is less likely to occur (Read et al., 2009; Schraiber et al., 2012). However, it has been shown that mosquitoes carrying life-shortening Wolbachia strains, such as *w*MelPop, have to be released in very high numbers compared to mosquito populations present at the release site, and that they cannot invade regional mosquito populations (Hughes & Britton, 2013; McMeniman et al., 2009; Walker et al., 2011). In contrast to the reduction of viral transmission due to Wolbachia-induced life-shortening, vector refractoriness is expected to diminish over time as a function of vector/Wolbachia co-adaptation (Vavre & Charlat, 2012).

As discussed previously, the density of *Wolbachia* in transinfected mosquitoes, and indeed in homologous mosquito/*Wolbachia* combinations, is crucial to the *Wolbachia*mediated antiviral activity (Lu *et al.*, 2012; Osborne *et al.*, 2009, 2012). Therefore, any decrease in *Wolbachia* density due to mosquito/*Wolbachia* co-adaptation may lead to a significant decrease in antiviral activity. However, it is clear from studies in *D. melanogaster* that long-term vector/ *Wolbachia*/virus associations can lead to continued protection against viruses. For example, invasion of *D. melanogaster* by *w*Mel occurred during the last 80 years (Riegler *et al.*, 2005), yet *w*Mel still confers protection against DCV (Teixeira *et al.*, 2008). This is encouraging and further studies in this model system could be extremely beneficial to the field.

If, at some point, reintroductions of *Wolbachia*-infected mosquitoes are required, then understanding the mechanism of CI becomes important. CI will confer an evolutionary advantage to released mosquitoes over the wild population, which aids the spread and maintenance of *Wolbachia* within the wild population. It will, however, also limit the spread of newly introduced *Wolbachia* strains into the same population, as CI will prevent successful mating of the incoming mosquitoes with individuals of the original population (see Fig. 1).

Further to CI, the introduction of novel mosquito/ *Wolbachia* combinations is hindered by the aforementioned technical difficulties of transinfections. As *Wolbachia* transinfection between closely related species is more successful (Bian *et al.*, 2010; Russell *et al.*, 2009), newly discovered *Wolbachia* strains of *Ae. bromeliae* (a YFV vector) and *Aedes metallicus* may be suitable candidates for introduction into aedine vectors (Osei-Poku *et al.*, 2012). Transinfection can also result in an evolutionary disadvantage within the mosquito, for example reduced fecundity or viability. This in turn could affect the spread within a population. Coupling *Wolbachia* introduction to insecticide resistance has been suggested in order to counteract these effects (Hoffmann & Turelli, 2013). Spread of such resistance genes into wild populations requires careful evaluation and is dependent on local mosquito populations and control strategies.

The impact of introduced *Wolbachia* on the target virus may be complex. Modelling indicates that different *Wolbachia* strains have varying abilities to reduce transmission frequencies (R_0) (Bian *et al.*, 2010; Hoffmann *et al.*, 2011; McMeniman *et al.*, 2009), and if transmission is only partially blocked *Wolbachia* will place the target virus in a bottleneck, perhaps leading to the evolution of a virus resistant to *Wolbachia*-mediated protection. Furthermore, modelling of the co-evolution of parasites and protecting symbionts suggests increases in parasite virulence (Jones *et al.*, 2011). The implication – that exposure to *Wolbachia* might result in increased virulence of DENV or CHIKV – is of obvious concern. There is clearly a need for long-term studies, perhaps in tissue culture, of the impact of *Wolbachia* on virus evolution.

Despite these complications, initial trials with transinfected mosquitoes are encouraging. Following cage experiments with *Ae. aegypti* and *w*AlbB (Xi *et al.*, 2005), experiments with *w*Mel and caged *Ae. aegypti* in semi-field conditions have shown rapid establishment of infection in the insect population and inhibition of DENV replication (Walker *et al.*, 2011). Release experiments into the Australian environment have achieved spread of *w*Mel into two natural *Ae. aegypti* populations within months of release of infected adult mosquitoes (Hoffmann *et al.*, 2011), with no detectable adverse effects on the environment and human health. Indeed, new field trials are set to begin in Australia (http://www.eliminatedengue.com/progress).

Wolbachia offers a unique opportunity to potentially understand and control arbovirus transmission. Indeed, more recent studies have indicated the use of *Wolbachia* to control *Plasmodium* infections, opening the field further (Bian *et al.*, 2013a). For virologists, investigating and potentially manipulating the mechanisms underlying the inhibition of virus transmission by *Wolbachia* will remain an interesting and challenging task.

Acknowledgements

Work in the authors' laboratory is funded by UK Medical Research Council, Wellcome Trust (094664/Z/10/Z) and the ERA-Net EMIDA (contract no. 219235). We thank William Sullivan, Laura Serbus and Alain Debec for sharing the Jw18 cells. We also thank Frazer Rixon for his help with electron microscopy.

References

Acosta, E. G., Castilla, V. & Damonte, E. B. (2008). Functional entry of dengue virus into *Aedes albopictus* mosquito cells is dependent on clathrin-mediated endocytosis. *J Gen Virol* **89**, 474–484.

Acosta, E. G., Castilla, V. & Damonte, E. B. (2009). Alternative infectious entry pathways for dengue virus serotypes into mammalian cells. *Cell Microbiol* **11**, 1533–1549.

Alphey, L., McKemey, A., Nimmo, D., Neira Oviedo, M., Lacroix, R., Matzen, K. & Beech, C. (2013). Genetic control of *Aedes* mosquitoes. *Pathog Glob Health* **107**, 170–179.

Armbruster, P., Damsky, W. E., Jr, Giordano, R., Birungi, J., Munstermann, L. E. & Conn, J. E. (2003). Infection of New- and Old-World *Aedes albopictus* (Diptera: Culicidae) by the intracellular parasite *Wolbachia*: implications for host mitochondrial DNA evolution. *J Med Entomol* **40**, 356–360.

Asgari, S. (2013). MicroRNA functions in insects. *Insect Biochem Mol Biol* 43, 388–397.

Avadhanula, V., Weasner, B. P., Hardy, G. G., Kumar, J. P. & Hardy, R. W. (2009). A novel system for the launch of alphavirus RNA synthesis reveals a role for the Imd pathway in arthropod antiviral response. *PLoS Pathog* 5, e1000582.

Bartholomay, L. C., Waterhouse, R. M., Mayhew, G. F., Campbell, C. L., Michel, K., Zou, Z., Ramirez, J. L., Das, S., Alvarez, K. & other authors (2010). Pathogenomics of *Culex quinquefasciatus* and metaanalysis of infection responses to diverse pathogens. *Science* 330, 88–90.

Berticat, C., Rousset, F., Raymond, M., Berthomieu, A. & Weill, M. (2002). High *Wolbachia* density in insecticide-resistant mosquitoes. *Proc Biol Sci* 269, 1413–1416.

Bian, G., Xu, Y., Lu, P., Xie, Y. & Xi, Z. (2010). The endosymbiotic bacterium *Wolbachia* induces resistance to dengue virus in *Aedes aegypti*. *PLoS Pathog* 6, e1000833.

Bian, G., Joshi, D., Dong, Y., Lu, P., Zhou, G., Pan, X., Xu, Y., Dimopoulos, G. & Xi, Z. (2013a). Wolbachia invades Anopheles stephensi populations and induces refractoriness to Plasmodium infection. Science 340, 748–751.

Bian, G., Zhou, G., Lu, P. & Xi, Z. (2013b). Replacing a native *Wolbachia* with a novel strain results in an increase in endosymbiont load and resistance to dengue virus in a mosquito vector. *PLoS Negl Trop Dis* 7, e2250.

Black, W. C., IV, Alphey, L. & James, A. A. (2011). Why RIDL is not SIT. *Trends Parasitol* 27, 362–370.

Blagrove, M. S., Arias-Goeta, C., Di Genua, C., Failloux, A. B. & Sinkins, S. P. (2013). A *Wolbachia w*Mel transinfection in *Aedes albopictus* is not detrimental to host fitness and inhibits chikungunya virus. *PLoS Negl Trop Dis* 7, e2152.

Blair, C. D. (2011). Mosquito RNAi is the major innate immune pathway controlling arbovirus infection and transmission. *Future Microbiol* **6**, 265–277.

Bourtzis, K., Pettigrew, M. M. & O'Neill, S. L. (2000). *Wolbachia* neither induces nor suppresses transcripts encoding antimicrobial peptides. *Insect Mol Biol* **9**, 635–639.

Brackney, D. E., Scott, J. C., Sagawa, F., Woodward, J. E., Miller, N. A., Schilkey, F. D., Mudge, J., Wilusz, J., Olson, K. E. & other authors (2010). C6/36 *Aedes albopictus* cells have a dysfunctional antiviral RNA interference response. *PLoS Negl Trop Dis* **4**, e856.

Brown, D. T. & Condreay, L. D. (1986). Replication of alphaviruses in mosquito cells. In *The Togaviridae and Flaviviridae*, pp. 171–207. Edited by M. J. Schlesinger. New York: Plenum Publishing.

Brownstein, J. S., Hett, E. & O'Neill, S. L. (2003). The potential of virulent *Wolbachia* to modulate disease transmission by insects. *J Invertebr Pathol* 84, 24–29.

Caragata, E. P., Rancès, E., Hedges, L. M., Gofton, A. W., Johnson, K. N., O'Neill, S. L. & McGraw, E. A. (2013). Dietary cholesterol modulates pathogen blocking by *Wolbachia*. *PLoS Pathog* 9, e1003459.

Carro, A. C. & Damonte, E. B. (2013). Requirement of cholesterol in the viral envelope for dengue virus infection. *Virus Res* 174, 78–87.

Chatterjee, P. K., Vashishtha, M. & Kielian, M. (2000). Biochemical consequences of a mutation that controls the cholesterol dependence of Semliki Forest virus fusion. *J Virol* **74**, 1623–1631.

Cho, K. O., Kim, G. W. & Lee, O. K. (2011). *Wolbachia* bacteria reside in host Golgi-related vesicles whose position is regulated by polarity proteins. *PLoS ONE* 6, e22703.

Christophides, G. K., Zdobnov, E., Barillas-Mury, C., Birney, E., Blandin, S., Blass, C., Brey, P. T., Collins, F. H., Danielli, A. & other authors (2002). Immunity-related genes and gene families in *Anopheles gambiae. Science* 298, 159–165.

Clayton, R. B. (1964). The utilization of sterols by insects. J Lipid Res 5, 3–19.

Costa, A., Jan, E., Sarnow, P. & Schneider, D. (2009). The Imd pathway is involved in antiviral immune responses in *Drosophila*. *PLoS ONE* **4**, e7436.

Dobson, S. L., Bourtzis, K., Braig, H. R., Jones, B. F., Zhou, W., Rousset, F. & O'Neill, S. L. (1999). *Wolbachia* infections are distributed throughout insect somatic and germ line tissues. *Insect Biochem Mol Biol* 29, 153–160.

Donald, C. L., Kohl, A. & Schnettler, E. (2012). New insights into control of arbovirus replication and spread by insect RNA interference pathways. *Insects* **3**, 511–531.

Dostert, C., Jouanguy, E., Irving, P., Troxler, L., Galiana-Arnoux, D., Hetru, C., Hoffmann, J. A. & Imler, J. L. (2005). The Jak-STAT signaling pathway is required but not sufficient for the antiviral response of *Drosophila*. *Nat Immunol* 6, 946–953.

Dutton, T. J. & Sinkins, S. P. (2004). Strain-specific quantification of *Wolbachia* density in *Aedes albopictus* and effects of larval rearing conditions. *Insect Mol Biol* 13, 317–322.

Echaubard, P., Duron, O., Agnew, P., Sidobre, C., Noël, V., Weill, M. & Michalakis, Y. (2010). Rapid evolution of *Wolbachia* density in insecticide resistant *Culex pipiens*. *Heredity* (*Edinb*) 104, 15–19.

Engelstädter, J. & Telschow, A. (2009). Cytoplasmic incompatibility and host population structure. *Heredity (Edinb)* 103, 196–207.

Fragkoudis, R., Chi, Y., Siu, R. W. C., Barry, G., Attarzadeh-Yazdi, G., Merits, A., Nash, A. A., Fazakerley, J. K. & Kohl, A. (2008). Semliki Forest virus strongly reduces mosquito host defence signaling. *Insect Mol Biol* 17, 647–656.

Fragkoudis, R., Attarzadeh-Yazdi, G., Nash, A. A., Fazakerley, J. K. & Kohl, A. (2009). Advances in dissecting mosquito innate immune responses to arbovirus infection. *J Gen Virol* **90**, 2061–2072.

Frentiu, F. D., Robinson, J., Young, P. R., McGraw, E. A. & O'Neill, S. L. (2010). *Wolbachia*-mediated resistance to dengue virus infection and death at the cellular level. *PLoS ONE* 5, e13398.

Glaser, R. L. & Meola, M. A. (2010). The native *Wolbachia* endosymbionts of *Drosophila melanogaster* and *Culex quinquefasciatus* increase host resistance to West Nile virus infection. *PLoS ONE* **5**, e11977.

Hafer, A., Whittlesey, R., Brown, D. T. & Hernandez, R. (2009). Differential incorporation of cholesterol by Sindbis virus grown in mammalian or insect cells. *J Virol* **83**, 9113–9121.

Haine, E. R., Pickup, N. J. & Cook, J. M. (2005). Horizontal transmission of *Wolbachia* in a *Drosophila* community. *Ecol Entomol* 30, 464–472.

Halstead, S. B. (2012). Dengue vaccine development: a 75 % solution? *Lancet* 380, 1535–1536.

Hancock, P. A. & Godfray, H. C. (2012). Modelling the spread of *Wolbachia* in spatially heterogeneous environments. *J R Soc Interface* 9, 3045–3054.

Heath, B. D., Butcher, R. D. J., Whitfield, W. G. F. & Hubbard, S. F. (1999). Horizontal transfer of *Wolbachia* between phylogenetically distant insect species by a naturally occurring mechanism. *Curr Biol* 9, 313–316.

Hedges, L. M., Brownlie, J. C., O'Neill, S. L. & Johnson, K. N. (2008). *Wolbachia* and virus protection in insects. *Science* **322**, 702.

Hedges, L. M., Yamada, R., O'Neill, S. L. & Johnson, K. N. (2012). The siRNA pathway is not essential for *Wolbachia*-mediated antiviral protection in *Drosophila*. *Appl Environ Microbiol* **78**, 6773–6776.

Hertig, M. & Wolbach, S. B. (1924). Studies on *Rickettsia*-like microorganisms in insects. J Med Res 44, 329–374, 7.

Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A. & Werren, J. H. (2008). How many species are infected with *Wolbachia*?–A statistical analysis of current data. *FEMS Microbiol Lett* 281, 215–220.

Hoffmann, A. A. & Turelli, M. (2013). Facilitating *Wolbachia* introductions into mosquito populations through insecticide-resistance selection. *Proc Biol Sci* 280, 20130371.

Hoffmann, A. A., Montgomery, B. L., Popovici, J., Iturbe-Ormaetxe, I., Johnson, P. H., Muzzi, F., Greenfield, M., Durkan, M., Leong, Y. S. & other authors (2011). Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. *Nature* 476, 454–457.

Huang, Z., Kingsolver, M. B., Avadhanula, V. & Hardy, R. W. (2013). An antiviral role for antimicrobial peptides during the arthropod response to alphavirus replication. *J Virol* 87, 4272–4280.

Hughes, H. & Britton, N. F. (2013). Modelling the use of *Wolbachia* to control dengue fever transmission. *Bull Math Biol* 75, 796–818.

Huigens, M. E., Luck, R. F., Klaassen, R. H. G., Maas, M. F. P. M., Timmermans, M. J. T. N. & Stouthamer, R. (2000). Infectious parthenogenesis. *Nature* 405, 178–179.

Huigens, M. E., de Almeida, R. P., Boons, P. A. H., Luck, R. F. & Stouthamer, R. (2004). Natural interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia* in *Trichogramma* wasps. *Proc Biol Sci* 271, 509–515.

Hussain, M., Frentiu, F. D., Moreira, L. A., O'Neill, S. L. & Asgari, S. (2011). *Wolbachia* uses host microRNAs to manipulate host gene expression and facilitate colonization of the dengue vector *Aedes aegypti*. *Proc Natl Acad Sci U S A* **108**, 9250–9255.

Hussain, M., Lu, G., Torres, S., Edmonds, J. H., Kay, B. H., Khromykh, A. A. & Asgari, S. (2013). Effect of *Wolbachia* on replication of West Nile virus in a mosquito cell line and adult mosquitoes. *J Virol* 87, 851–858.

Jones, E. O., White, A. & Boots, M. (2011). The evolution of host protection by vertically transmitted parasites. *Proc Biol Sci* 278, 863–870.

Kambhampati, S., Rai, K. S. & Burgun, S. J. (1993). Unidirectional cytoplasmic incompatibility in the mosquito, *Aedes-Albopictus*. *Evolution* **47**, 673–677.

Kambris, Z., Blagborough, A. M., Pinto, S. B., Blagrove, M. S., Godfray, H. C., Sinden, R. E. & Sinkins, S. P. (2010). *Wolbachia* stimulates immune gene expression and inhibits plasmodium development in *Anopheles gambiae*. *PLoS Pathog* 6, e1001143.

Kielian, M. (1995). Membrane fusion and the alphavirus life cycle. *Adv Virus Res* 45, 113–151.

Kittayapong, P., Baisley, K. J., Baimai, V. & O'Neill, S. L. (2000). Distribution and diversity of *Wolbachia* infections in Southeast Asian mosquitoes (Diptera: Culicidae). *J Med Entomol* **37**, 340–345.

Kraaijeveld, K., Franco, P., de Knijff, P., Stouthamer, R. & van Alphen, J. J. (2011). Clonal genetic variation in a *Wolbachia*-infected asexual wasp: horizontal transmission or historical sex? *Mol Ecol* 20, 3644–3652.

Krejbich-Trotot, P., Gay, B., Li-Pat-Yuen, G., Hoarau, J. J., Jaffar-Bandjee, M. C., Briant, L., Gasque, P. & Denizot, M. (2011). Chikungunya triggers an autophagic process which promotes viral replication. *Virol J* 8, 432.

Kremer, N., Voronin, D., Charif, D., Mavingui, P., Mollereau, B. & Vavre, F. (2009). *Wolbachia* interferes with ferritin expression and iron metabolism in insects. *PLoS Pathog* 5, e1000630.

Lacroix, R., McKemey, A. R., Raduan, N., Kwee Wee, L., Hong Ming, W., Guat Ney, T., Rahidah A A, S., Salman, S., Subramaniam, S. & other authors (2012). Open field release of genetically engineered sterile male *Aedes aegypti* in Malaysia. *PLoS ONE* 7, e42771.

Landmann, F., Orsi, G. A., Loppin, B. & Sullivan, W. (2009). *Wolbachia*-mediated cytoplasmic incompatibility is associated with impaired histone deposition in the male pronucleus. *PLoS Pathog* 5, e1000343.

Lange, Y., Ye, J., Rigney, M. & Steck, T. L. (1999). Regulation of endoplasmic reticulum cholesterol by plasma membrane cholesterol. *J Lipid Res* 40, 2264–2270.

Laven, H. (1967). Eradication of *Culex pipiens fatigans* through cytoplasmic incompatibility. *Nature* 216, 383–384.

Le Clec'h, W., Braquart-Varnier, C., Raimond, M., Ferdy, J. B., Bouchon, D. & Sicard, M. (2012). High virulence of *Wolbachia* after host switching: when autophagy hurts. *PLoS Pathog* 8, e1002844.

Lee, Y. R., Lei, H. Y., Liu, M. T., Wang, J. R., Chen, S. H., Jiang-Shieh, Y. F., Lin, Y. S., Yeh, T. M., Liu, C. C. & Liu, H. S. (2008). Autophagic machinery activated by dengue virus enhances virus replication. *Virology* **374**, 240–248.

Lemaitre, B. & Hoffmann, J. (2007). The host defense of Drosophila melanogaster. Annu Rev Immunol 25, 697–743.

Lin, M. & Rikihisa, Y. (2003). *Ehrlichia chaffeensis* and *Anaplasma phagocytophilum* lack genes for lipid A biosynthesis and incorporate cholesterol for their survival. *Infect Immun* 71, 5324–5331.

Lipsitch, M. & Moxon, E. R. (1997). Virulence and transmissibility of pathogens: what is the relationship? *Trends Microbiol* 5, 31–37.

Lu, Y. E., Cassese, T. & Kielian, M. (1999). The cholesterol requirement for Sindbis virus entry and exit and characterization of a spike protein region involved in cholesterol dependence. *J Virol* 73, 4272–4278.

Lu, P., Bian, G., Pan, X. & Xi, Z. (2012). *Wolbachia* induces densitydependent inhibition to dengue virus in mosquito cells. *PLoS Negl Trop Dis* 6, e1754.

Mackenzie, J. M., Khromykh, A. A. & Parton, R. G. (2007). Cholesterol manipulation by West Nile virus perturbs the cellular immune response. *Cell Host Microbe* **2**, 229–239.

Marquardt, M. T., Phalen, T. & Kielian, M. (1993). Cholesterol is required in the exit pathway of Semliki Forest virus. *J Cell Biol* 123, 57–65.

McMeniman, C. J., Lane, A. M., Fong, A. W., Voronin, D. A., Iturbe-Ormaetxe, I., Yamada, R., McGraw, E. A. & O'Neill, S. L. (2008). Host adaptation of a *Wolbachia* strain after long-term serial passage in mosquito cell lines. *Appl Environ Microbiol* 74, 6963–6969.

McMeniman, C. J., Lane, R. V., Cass, B. N., Fong, A. W., Sidhu, M., Wang, Y. F. & O'Neill, S. L. (2009). Stable introduction of a lifeshortening *Wolbachia* infection into the mosquito *Aedes aegypti*. *Science* 323, 141–144.

Merkling, S. H. & van Rij, R. P. (2013). Beyond RNAi: antiviral defense strategies in *Drosophila* and mosquito. J Insect Physiol 59, 159–170.

Miller, W. J. & Riegler, M. (2006). Evolutionary dynamics of wAu-like *Wolbachia* variants in neotropical *Drosophila* spp. *Appl Environ Microbiol* 72, 826–835.

Min, K. T. & Benzer, S. (1997). *Wolbachia*, normally a symbiont of *Drosophila*, can be virulent, causing degeneration and early death. *Proc Natl Acad Sci U S A* 94, 10792–10796.

Mitsuhashi, J., Nakasone, S. & Horie, Y. (1983). Sterol-free eukaryotic cells from continuous cell lines of insects. *Cell Biol Int Rep* 7, 1057–1062.

Moreira, L. A., Iturbe-Ormaetxe, I., Jeffery, J. A., Lu, G., Pyke, A. T., Hedges, L. M., Rocha, B. C., Hall-Mendelin, S., Day, A. & other authors (2009). A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, chikungunya, and *Plasmodium. Cell* 139, 1268–1278.

Mosso, C., Galván-Mendoza, I. J., Ludert, J. E. & del Angel, R. M. (2008). Endocytic pathway followed by dengue virus to infect the mosquito cell line C6/36 HT. *Virology* **378**, 193–199.

Mousson, L., Martin, E., Zouache, K., Madec, Y., Mavingui, P. & Failloux, A. B. (2010). *Wolbachia* modulates chikungunya replication in *Aedes albopictus. Mol Ecol* **19**, 1953–1964.

Mousson, L., Zouache, K., Arias-Goeta, C., Raquin, V., Mavingui, P. & Failloux, A. B. (2012). The native *Wolbachia* symbionts limit transmission of dengue virus in *Aedes albopictus*. *PLoS Negl Trop Dis* **6**, e1989.

Osborne, S. E., Leong, Y. S., O'Neill, S. L. & Johnson, K. N. (2009). Variation in antiviral protection mediated by different *Wolbachia* strains in *Drosophila simulans*. *PLoS Pathog* 5, e1000656.

Osborne, S. E., Iturbe-Ormaetxe, I., Brownlie, J. C., O'Neill, S. L. & Johnson, K. N. (2012). Antiviral protection and the importance of *Wolbachia* density and tissue tropism in *Drosophila simulans*. Appl Environ Microbiol **78**, 6922–6929.

Osei-Amo, S., Hussain, M., O'Neill, S. L. & Asgari, S. (2012). *Wolbachia*-induced aae-miR-12 miRNA negatively regulates the expression of MCT1 and MCM6 genes in *Wolbachia*-infected mosquito cell line. *PLoS ONE* 7, e50049.

Osei-Poku, J., Han, C., Mbogo, C. M. & Jiggins, F. M. (2012). Identification of *Wolbachia* strains in mosquito disease vectors. *PLoS ONE* **7**, e49922.

Pan, X., Zhou, G., Wu, J., Bian, G., Lu, P., Raikhel, A. S. & Xi, Z. (2012). Wolbachia induces reactive oxygen species (ROS)-dependent activation of the Toll pathway to control dengue virus in the mosquito *Aedes* aegypti. *Proc Natl Acad Sci U S A* **109**, E23–E31.

Phalen, T. & Kielian, M. (1991). Cholesterol is required for infection by Semliki Forest virus. J Cell Biol 112, 615–623.

Popovici, J., Moreira, L. A., Poinsignon, A., Iturbe-Ormaetxe, I., McNaughton, D. & O'Neill, S. L. (2010). Assessing key safety concerns of a *Wolbachia*-based strategy to control dengue transmission by *Aedes* mosquitoes. *Mem Inst Oswaldo Cruz* 105, 957–964.

Rancès, E., Ye, Y. H., Woolfit, M., McGraw, E. A. & O'Neill, S. L. (2012). The relative importance of innate immune priming in *Wolbachia*-mediated dengue interference. *PLoS Pathog* 8, e1002548.

Rancès, E., Johnson, T. K., Popovici, J., Iturbe-Ormaetxe, I., Zakir, T., Warr, C. G. & O'Neill, S. L. (2013). The toll and Imd pathways are not required for *Wolbachia*-mediated dengue virus interference. *J Virol* 87, 11945–11949.

Read, A. F., Lynch, P. A. & Thomas, M. B. (2009). How to make evolution-proof insecticides for malaria control. *PLoS Biol* 7, e1000058.

Riegler, M., Sidhu, M., Miller, W. J. & O'Neill, S. L. (2005). Evidence for a global *Wolbachia* replacement in *Drosophila melanogaster. Curr Biol* **15**, 1428–1433.

Rodriguez-Andres, J., Rani, S., Varjak, M., Chase-Topping, M. E., Beck, M. H., Ferguson, M. C., Schnettler, E., Fragkoudis, R., Barry, G. & other authors (2012). Phenoloxidase activity acts as a mosquito innate immune response against infection with Semliki Forest virus. *PLoS Pathog* **8**, e1002977.

Rothwell, C., Lebreton, A., Young Ng, C., Lim, J. Y., Liu, W., Vasudevan, S., Labow, M., Gu, F. & Gaither, L. A. (2009). Cholesterol biosynthesis modulation regulates dengue viral replication. *Virology* **389**, 8–19.

Rottschaefer, S. M. & Lazzaro, B. P. (2012). No effect of *Wolbachia* on resistance to intracellular infection by pathogenic bacteria in *Drosophila melanogaster*. *PLoS ONE* 7, e40500.

Russell, J. A., Goldman-Huertas, B., Moreau, C. S., Baldo, L., Stahlhut, J. K., Werren, J. H. & Pierce, N. E. (2009). Specialization and geographic isolation among *Wolbachia* symbionts from ants and lycaenid butterflies. *Evolution* **63**, 624–640.

Schofield, P. (2002). Spatially explicit models of Turelli-Hoffmann *Wolbachia* invasive wave fronts. *J Theor Biol* **215**, 121–131.

Schraiber, J. G., Kaczmarczyk, A. N., Kwok, R., Park, M., Silverstein, R., Rutaganira, F. U., Aggarwal, T., Schwemmer, M. A., Hom, C. L. & other authors (2012). Constraints on the use of lifespan-shortening *Wolbachia* to control dengue fever. *J Theor Biol* 297, 26–32.

Scott, J. C., Brackney, D. E., Campbell, C. L., Bondu-Hawkins, V., Hjelle, B., Ebel, G. D., Olson, K. E. & Blair, C. D. (2010). Comparison of dengue virus type 2-specific small RNAs from RNA interferencecompetent and -incompetent mosquito cells. *PLoS Negl Trop Dis* 4, e848.

Serbus, L. R., Casper-Lindley, C., Landmann, F. & Sullivan, W. (2008). The genetics and cell biology of *Wolbachia*-host interactions. *Annu Rev Genet* **42**, 683–707.

Serbus, L. R., Landmann, F., Bray, W. M., White, P. M., Ruybal, J., Lokey, R. S., Debec, A. & Sullivan, W. (2012). A cell-based screen reveals that the albendazole metabolite, albendazole sulfone, targets *Wolbachia. PLoS Pathog* 8, e1002922.

Shaw, A. E., Veronesi, E., Maurin, G., Ftaich, N., Guiguen, F., Rixon, F., Ratinier, M., Mertens, P., Carpenter, S. & other authors (2012). *Drosophila melanogaster* as a model organism for bluetongue virus replication and tropism. *J Virol* 86, 9015–9024.

Silberkang, M., Havel, C. M., Friend, D. S., McCarthy, B. J. & Watson, J. A. (1983). Isoprene synthesis in isolated embryonic *Drosophila* cells. I. Sterol-deficient eukaryotic cells. *J Biol Chem* 258, 8503–8511.

Sinkins, S. P. (2004). *Wolbachia* and cytoplasmic incompatibility in mosquitoes. *Insect Biochem Mol Biol* 34, 723–729.

Sinkins, S. P., Braig, H. R. & O'Neill, S. L. (1995). *Wolbachia* superinfections and the expression of cytoplasmic incompatibility. *Proc Biol Sci* 261, 325–330.

Smartt, C. T., Richards, S. L., Anderson, S. L. & Erickson, J. S. (2009). West Nile virus infection alters midgut gene expression in *Culex pipiens quinquefasciatus* Say (Diptera: Culicidae). *Am J Trop Med Hyg* **81**, 258–263.

Smit, J. M., Bittman, R. & Wilschut, J. (1999). Low-pH-dependent fusion of Sindbis virus with receptor-free cholesterol- and sphingo-lipid-containing liposomes. *J Virol* 73, 8476–8484.

Sousa, I. P., Jr, Carvalho, C. A., Ferreira, D. F., Weissmüller, G., Rocha, G. M., Silva, J. L. & Gomes, A. M. (2011). Envelope lipid-packing as a critical factor for the biological activity and stability of alphavirus particles isolated from mammalian and mosquito cells. *J Biol Chem* 286, 1730–1736.

Souza-Neto, J. A., Sim, S. & Dimopoulos, G. (2009). An evolutionary conserved function of the JAK-STAT pathway in anti-dengue defense. *Proc Natl Acad Sci U S A* 106, 17841–17846.

Stouthamer, R., Breeuwer, J. A. J. & Hurst, G. D. D. (1999). Wolbachia pipientis: microbial manipulator of arthropod reproduction. Annu Rev Microbiol 53, 71–102.

Tchankouo-Nguetcheu, S., Khun, H., Pincet, L., Roux, P., Bahut, M., Huerre, M., Guette, C. & Choumet, V. (2010). Differential protein modulation in midguts of *Aedes aegypti* infected with chikungunya and dengue 2 viruses. *PLoS ONE* 5, e13149.

Teixeira, L., Ferreira, A. & Ashburner, M. (2008). The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol* 6, e2.

Thomas, P., Kenny, N., Eyles, D., Moreira, L. A., O'Neill, S. L. & Asgari, S. (2011). Infection with the wMel and wMelPop strains of *Wolbachia* leads to higher levels of melanization in the hemolymph of *Drosophila* melanogaster, *Drosophila simulans* and *Aedes aegypti. Dev Comp Immunol* **35**, 360–365.

Tram, U. & Sullivan, W. (2002). Role of delayed nuclear envelope breakdown and mitosis in *Wolbachia*-induced cytoplasmic incompatibility. *Science* 296, 1124–1126.

Tram, U., Fredrick, K., Werren, J. H. & Sullivan, W. (2006). Paternal chromosome segregation during the first mitotic division determines *Wolbachia*-induced cytoplasmic incompatibility phenotype. *J Cell Sci* 119, 3655–3663.

Tsai, K. H., Huang, C. G., Wu, W. J., Chuang, C. K., Lin, C. C. & Chen, W. J. (2006). Parallel infection of Japanese encephalitis virus and *Wolbachia* within cells of mosquito salivary glands. *J Med Entomol* **43**, 752–756.

Tsetsarkin, K. A., McGee, C. E. & Higgs, S. (2011). Chikungunya virus adaptation to *Aedes albopictus* mosquitoes does not correlate with acquisition of cholesterol dependence or decreased pH threshold for fusion reaction. *Virol J* **8**, 376.

Turelli, M. (2010). Cytoplasmic incompatibility in populations with overlapping generations. *Evolution* 64, 232–241.

Turelli, M. & Hoffmann, A. A. (1991). Rapid spread of an inherited incompatibility factor in California *Drosophila*. *Nature* 353, 440–442.

Turelli, M. & Hoffmann, A. A. (1995). Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations. *Genetics* 140, 1319–1338.

Umashankar, M., Sánchez-San Martín, C., Liao, M., Reilly, B., Guo, A., Taylor, G. & Kielian, M. (2008). Differential cholesterol binding by class II fusion proteins determines membrane fusion properties. *J Virol* 82, 9245–9253.

van den Hurk, A. F., Hall-Mendelin, S., Pyke, A. T., Frentiu, F. D., McElroy, K., Day, A., Higgs, S. & O'Neill, S. L. (2012). Impact of *Wolbachia* on infection with chikungunya and yellow fever viruses in the mosquito vector *Aedes aegypti*. *PLoS Negl Trop Dis* 6, e1892.

Vavre, F. & Charlat, S. (2012). Making (good) use of *Wolbachia*: what the models say. *Curr Opin Microbiol* 15, 263–268.

Vavre, F., Fleury, F., Lepetit, D., Fouillet, P. & Boulétreau, M. (1999). Phylogenetic evidence for horizontal transmission of *Wolbachia* in host-parasitoid associations. *Mol Biol Evol* 16, 1711–1723.

Voronin, D., Cook, D. A., Steven, A. & Taylor, M. J. (2012). Autophagy regulates *Wolbachia* populations across diverse symbiotic associations. *Proc Natl Acad Sci U S A* 109, E1638–E1646.

Waldock, J., Olson, K. E. & Christophides, G. K. (2012). Anopheles gambiae antiviral immune response to systemic o'nyong-nyong infection. *PLoS Negl Trop Dis* 6, e1565.

Walker, T., Johnson, P. H., Moreira, L. A., Iturbe-Ormaetxe, I., Frentiu, F. D., McMeniman, C. J., Leong, Y. S., Dong, Y., Axford, J. & other authors (2011). The wMel *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti* populations. *Nature* 476, 450–453.

Waterhouse, R. M., Kriventseva, E. V., Meister, S., Xi, Z., Alvarez, K. S., Bartholomay, L. C., Barillas-Mury, C., Bian, G., Blandin, S. & other authors (2007). Evolutionary dynamics of immune-related genes and pathways in disease-vector mosquitoes. *Science* **316**, 1738–1743. Weaver, S. C. & Reisen, W. K. (2010). Present and future arboviral threats. *Antiviral Res* 85, 328–345.

Welsch, S., Miller, S., Romero-Brey, I., Merz, A., Bleck, C. K., Walther, P., Fuller, S. D., Antony, C., Krijnse-Locker, J. & Bartenschlager, R. (2009). Composition and three-dimensional architecture of the dengue virus replication and assembly sites. *Cell Host Microbe* 5, 365–375.

Werren, J. H., Zhang, W. & Guo, L. R. (1995). Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. *Proc Biol Sci* 261, 55–63.

Werren, J. H., Baldo, L. & Clark, M. E. (2008). Wolbachia: master manipulators of invertebrate biology. Nat Rev Microbiol 6, 741–751.

Wilke, A. B. & Marrelli, M. T. (2012). Genetic control of mosquitoes: population suppression strategies. *Rev Inst Med Trop Sao Paulo* 54, 287–292.

Wong, Z. S., Hedges, L. M., Brownlie, J. C. & Johnson, K. N. (2011). *Wolbachia*-mediated antibacterial protection and immune gene regulation in Drosophila. *PLoS ONE* 6, e25430.

Xi, Z., Khoo, C. C. & Dobson, S. L. (2005). *Wolbachia* establishment and invasion in an Aedes aegypti laboratory population. *Science* 310, 326–328.

Xi, Z., Ramirez, J. L. & Dimopoulos, G. (2008). The Aedes aegypti toll pathway controls dengue virus infection. *PLoS Pathog* 4, e1000098.

Ye, Y. H., Woolfit, M., Huttley, G. A., Rances, E., Caragata, E. P., Popovici, J., O'Neill, S. L. & McGraw, E. A. (2013). Infection with a virulent strain of *Wolbachia* disrupts genome wide-patterns of cytosine methylation in the mosquito *Aedes aegypti*. *PLoS ONE* **8**, e66482.

Yeap, H. L., Mee, P., Walker, T., Weeks, A. R., O'Neill, S. L., Johnson, P., Ritchie, S. A., Richardson, K. M., Doig, C. & other authors (2011). Dynamics of the "popcorn" *Wolbachia* infection in outbred *Aedes aegypti* informs prospects for mosquito vector control. *Genetics* 187, 583–595.

Yen, J. H. & Barr, A. R. (1971). New hypothesis of the cause of cytoplasmic incompatibility in *Culex pipiens* L. *Nature* 232, 657–658.

Yen, J. H. & Barr, A. R. (1973). The etiological agent of cytoplasmic incompatibility in *Culex pipiens*. J Invertebr Pathol 22, 242–250.

Zambon, R. A., Nandakumar, M., Vakharia, V. N. & Wu, L. P. (2005). The Toll pathway is important for an antiviral response in *Drosophila*. *Proc Natl Acad Sci U S A* **102**, 7257–7262.

Zhang, G., Hussain, M., O'Neill, S. L. & Asgari, S. (2013). *Wolbachia* uses a host microRNA to regulate transcripts of a methyltransferase, contributing to dengue virus inhibition in *Aedes aegypti. Proc Natl Acad Sci U S A* **110**, 10276–10281.

Zug, R. & Hammerstein, P. (2012). Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected. *PLoS ONE* 7, e38544.